

The routine childhood immunisation schedule for Wales

from 1 October 2026

Age due	Diseases protected against		Vaccine and name		Usual site
8 weeks old	Diphtheria, tetanus, pertussis (whooping cough), polio, Haemophilus influenzae type b (Hib) and hepatitis B		DTaP/IPV/Hib/HepB (6-in-1)	Infanrix hexa or Vaxelis	Thigh
	Meningococcal group B (MenB)		MenB	Bexsero	Thigh
	Rotavirus gastroenteritis		Rotavirus	Rotarix	By mouth
12 weeks old	Diphtheria, tetanus, pertussis, polio, Hib and hepatitis B		DTaP/IPV/Hib/HepB (6-in-1)	Infanrix hexa or Vaxelis	Thigh
	MenB		MenB	Bexero	Thigh
	Rotavirus gastroenteritis		Rotavirus	Rotarix	By mouth
16 weeks old	Diphtheria, tetanus, pertussis, polio, Hib and hepatitis B		DTaP/IPV/Hib/HepB (6-in-1)	Infanrix hexa or Vaxelis	Thigh
	Pneumococcal		PCV13	Prevenar 13 ¹	Thigh
12 months old	Pneumococcal		PCV13	Prevenar 13 ¹	Upper arm/ thigh
	MenB		MenB	Bexsero	Thigh
	Measles, mumps, rubella and varicella (chickenpox)		MMRV	ProQuad or Priorix-Tetra	Upper arm/ thigh
18 months old	Born before 1 July 2024	Born on or after 1 July 2024			
	No appointment	Measles, mumps, rubella and varicella (chickenpox) Diphtheria, tetanus, pertussis, polio, Hib and hepatitis B	MMRV DTaP/IPV/Hib/ HepB (6-in-1)	ProQuad or Priorix-Tetra Infanrix hexa or Vaxelis	Upper arm/ thigh
2 ² and 3 years old and all school aged children	Influenza (annually from September)		Live attenuated influenza vaccine	Fluenz ³	Both nostrils
3 years 4 months old	Born before 1 January 2025	Born on or after 1 January 2025			
	Diphtheria, tetanus, pertussis and polio Measles, mumps, rubella and varicella (chickenpox)	Diphtheria, tetanus, pertussis and polio No MMRV vaccine	dTaP/IPV (4-in-1) MMRV	Repevax ProQuad or Priorix-Tetra	Upper arm
School year 8 (12 to 13 year olds)	Cancers and genital warts caused by specific human papillomavirus (HPV) types		HPV ⁴ (one dose)	Gardasil 9	Upper arm
School year 9 (13 to 14 year olds)	Tetanus, diphtheria and polio		Td/IPV (3-in-1) (check MMR status)	Revaxis	Upper arm
	Meningococcal groups A, C, W and Y		MenACWY	Nimenrix or MenQuadfi	Upper arm

¹ Refer to the [Green Book Chapter: Pneumococcal - table 3 \(external site\)](#) for babies with asplenia, splenic dysfunction, complement disorder or severely immunocompromised.

² Children must be 2 years old by 31 August to receive influenza vaccine in the routine programme in autumn/winter.

³ If Fluenz is unsuitable, use inactivated flu vaccine.

⁴ Check the relevant chapter of the Green Book for individuals requiring a 3-dose schedule.

Selective childhood immunisation programmes¹

Target group	Age and schedule	Disease	Vaccines
Babies born to hepatitis B infected mothers	At birth and 4 weeks of age ²	Hepatitis B	Hepatitis B vaccines Engerix B or HBvaxPRO
Infants in areas of the country with TB incidence $\geq 40/100,000$	From birth	Tuberculosis	BCG
Infants with a parent or grandparent born in a high incidence country ³	From birth	Tuberculosis	BCG
Children in a risk group for influenza ⁴	From 6 months to 17 years	Influenza	Age-appropriate flu vaccine ⁴

¹ Other vaccines may be recommended for individuals with certain underlying medical conditions

² In addition to the hexavalent vaccine (Infanrix hexa or Vaxelis) given at 8, 12, 16 weeks and, for children born on or after 1 July 2024, also at 18 months. Take a blood sample for HBsAg at 12 to 18 months to exclude infection. See the [Selective neonatal Hepatitis B immunisation pathway | GOV.WALES \(external site\)](#)

³ Where the annual incidence of TB is $\geq 40/100,000$ see the [WHO country profile \(external site\)](#)

⁴ Refer to annual flu [Welsh Health Circular \(external site\)](#) for eligibility criteria and vaccine recommendations.

Yr amserlen imiwneiddio reolaidd i blant ar gyfer Cymru

o fis Hydref 2026

Yr oedran i'w roi	Yr afiechydon mae'n gwarchod rhag eu cael		Brechiad ac enw		Lleoliad arferol
8 wythnos oed	Difftheria, tetanws, pertwsis (y pas), polio, Haemophilus influenzae math b (Hib) a hepatitis B		DTaP/IPV/Hib/HepB (6-mewn-1)	Infanrix hexa® neu Vaxelis	Clun
	Grŵp meningococol B (MenB)		MenB	Bexsero	Clun
	Rotafeirws gastroenteritis		Rotafeirws	Rotarix	Ceg
12 wythnos oed	Difftheria, tetanws, pertwsis, polio, Hib a hepatitis B		DTaP/IPV/Hib/HepB (6-mewn-1)	Infanrix hexa neu Vaxelis	Clun
	MenB		MenB	Bexero	Clun
	Rotafeirws gastroenteritis		Rotafeirws	Rotarix	Ceg
16 wythnos oed	Difftheria, tetanws, pertwsis, polio, Hib a hepatitis B		DTaP/IPV/Hib/HepB (6-mewn-1)	Infanrix hexa® neu Vaxelis	Clun
	Niwmococol		PCV13	Prevenar 13 ¹	Clun
12 mis oed	Niwmococol		PCV13	Prevenar 13 ¹	Rhan uchaf y fraich/clun
	MenB		MenB	Bexsero	Clun
	Y frech goch, clwy'r pennau, rwbela a farisela (brech yr ieir)		MMRV	ProQuad neu Priorix-Tetra	Rhan uchaf y fraich/ clun
18 mis oed	Ganwyd cyn 1 Gorffennaf 2024	Ganwyd ar neu ar ôl 1 Gorffennaf 2024			
	Dim apwyntiad	Y frech goch, clwy'r pennau, rwbela a farisela (brech yr ieir) Difftheria, tetanws, pertwsis, polio, Hib a hepatitis B	MMRV DTaP/IPV/Hib/ HepB (6-mewn-1)	ProQuad neu Priorix-Tetra Infanrix hexa neu Vaxelis	Rhan uchaf y fraich/ clun
2 ² a 3 oed a phob plentyn oedran ysgol	Y Ffliw (bob blwyddyn o fis Medi)		Brechlyn fflw gwanedig byw (LAIV)	Fluenz ³	Dau dwll y trwyn
3 blynedd a 4 mis oed	Ganwyd cyn 1 Ionawr 2025	Ganwyd ar neu ar ôl 1 Ionawr 2025			
	Difftheria, tetanws, pertwsis a pholio Y frech goch, clwy'r pennau, rwbela a farisela (brech yr ieir)	Difftheria, tetanws, pertwsis a pholio Dim brechlyn MMRV	dTaP/IPV (4-mewn-1) MMRV	Repevax ProQuad neu Priorix-Tetra	Rhan uchaf y fraich
Blwyddyn ysgol 8 (12 i 13 oed)	Canserau a dafadennau gwenerol a achosir gan fathau penodol o'r feirws papiloma dynol (HPV)		HPV ⁴ (un dos)	Gardasil 9	Rhan uchaf y fraich
Blwyddyn ysgol 9 (13 a 14 oed)	Tetanws, difftheria a pholio		Td/IPV (3-mewn-1) (gwirio statws MMR)	Revaxis	Rhan uchaf y fraich
	Grwpiau meningococol A, C, W ac Y		MenACWY	Nimenrix neu MenQuadfi	Rhan uchaf y fraich

¹ Cyfeiriwch at [Bennod y Llyfr Gwyrdd: Niwmococol, tabl 3 \(safle allanol, Saesneg yn unig\)](#), ar gyfer babanod ag asplenia, camweithrediad y dduwg, anhwylder cyflenwol neu imiwnedd gwan iawn.

² Rhaid i blant fod yn 2 oed erbyn 31 Awst i gael brechlyn fflw fel rhan o'r rhaglen reolaidd yn yr hydref/gaeaf.

³ Os nad yw Fluenz® yn addas, defnyddiwch frechlyn fflw wedi'i anactifadu.

⁴ Gwiriwch y bennod berthnasol yn y Llyfr Gwyrdd os oes gennych unigolion sydd angen amserlen 3 dos.

Rhaglenni imiwneiddio dethol ar gyfer plant¹

Grwp targed	Oedran ac amserlen	Afiechyd	Brechciadau
Babanod wedi'u geni i famau wedi'u heintio â hepatitis B	Ar enedigaeth ac yn 4 wythnos oed ²	Hepatitis B	Brechciadau Hepatitis B Engerix B neu HBvaxPRO
Babanod mewn ardaloedd o'r wlad lle mae achosion o TB >= 40/100,000	O enedigaeth	Twbercwlosis	BCG
Babanod gyda rhiant neu daid neu nain wedi'u geni mewn gwlad gyda llawer o achosion ³	O enedigaeth	Twbercwlosis	BCG
Plant mewn grŵp risg ar gyfer y fflw ⁴	Rhwng 6 mis a 17 oed	Y fflw	Brechlyn fflw sy'n briodol i oedran ⁴

¹ Efallai y caiff brechciadau eraill eu hargymell ar gyfer unigolion gyda rhai cyflyrau.

² Yn ogystal â'r brechlyn hexavalent (Infanrix hexa neu Vaxelis) a roddir yn 8, 12, 16 wythnos ac, i blant a anwyd ar neu ar ôl 1 Gorffennaf 2024, hefyd yn 18 mis oed. Cymerwch sampl gwaed ar gyfer HBsAg rhwng 12 a 18 mis oed i eithrio haint. Gweler y [Llyfr imiwneiddio detholus newyddenedigol ar gyfer Hepatitis B | LLYWODRAETH CYMRU \(safle allanol\)](#)

³ Pan fo nifer yr achosion blynyddol o TB o leiaf 40 fesul 100,000, gweler [Proffil gwlad TB WHO \(Saesneg yn unig\)](#)

⁴ Cyfeiriwch at y fflw blynyddol [Cylchlythyrau iechyd \(safle allanol\)](#) am feini prawf cymhwystara ac argymhellion am frechlynnau