

lechyd Cyhoeddus Cymru Public Health Wales



Orthopaedic Surgical Site Infection Surveillance

2018 Annual Report:

All Wales

Includes data from 01/01/2008 – 31/12/2018

Version 1 Issued: 29/05/2019 The Healthcare Associated Infection and Antimicrobial Resistance Programme can be accessed via the Public Health Wales website at: http://www.publichealthwales.org

Published by Public Health Wales NHS Trust, No. 2 Capital Quarter, Tyndall Street, Cardiff CF10 4BZ

Published 29 May 2019, Version 1

Acknowledgements

We are grateful to the surgical teams, infection control teams and audit teams from all participating hospitals for continuing to provide surveillance data to Public Health Wales.

Report authors: Daryn Sutton, Wendy Harrison

Reference this document as:

Public Health Wales NHS Trust. 2018 Annual All Wales report on Orthopaedic Surgical Site Infection 2018, Wales.

Data requests and queries should be addressed to:

Public Health Wales NHS Trust HCAI and AMR Programme 4th floor, No. 2 Capital Quarter Tyndall Street Cardiff CF10 4BZ

Email: <u>harp@wales.nhs.uk</u>

Table of Contents

Tables2)
Figures	<u>)</u>
Introduction	3
Data interpretation	3
Summary	ł
Section 1: SSI rates	5
Section 2: Frequency of revisions	5
Annual Revisions6	5
Section 2: Procedure details and demographics7	7
Procedure types)
Prophylactic antibiotics)
Section 3: Post-procedure details11	Ĺ
Microbiology results	Ĺ
Section 4: Data quality)
Discussion	3
References	ł
Appendix15	5

Tables

Table 1 - SSI rates split by procedure type and demographics	7
Table 2 - Procedure type by OPCS code	9
Table 3 - Prophylactic antibiotics SSI rate	10
Table 4 - Microbiological results, organisms and SSI rates	11
Table 5 - Healthboard level SSI rates for 2018	15
Table 6 - Healthboard microbiological data for 2018	15

Figures

Figure 1 - Graph for All Wales primary hip SSI rate 2008-2018	5
Figure 2 - Graph for All Wales primary knee SSI rate 2008-2018	5
Figure 3 - Graph showing hip revision frequency in all Wales 2008-2018	6
Figure 4 - Graph showing knee revision frequency in all Wales 2008-2018	6
Figure 5 - Graph showing frequency of different procedure types 2008-2018	8
Figure 6 - Graph showing the usage of prophylactic antibiotics in 2017 vs 2018	10
Figure 7 - Proportions of data reporting sources per year 2008-2018	12

Introduction

The Healthcare Associated Infections team at Public Health Wales were instructed by the Welsh Government to develop and support the implementation of surveillance following orthopaedic procedures undertaken in NHS hospitals in Wales. This process has been mandatory since 2007.

Surgical Site Infection (SSI) is an important area for surveillance and remains a complication of surgery where human and financial costs are high [1] [2]. Additionally, most infections are preventable [3].

Serious patient consequences can result from SSIs, including pain, suffering and, on some occasions, they require additional surgical intervention [4]. It is important to recognise that SSIs can range from relatively trivial wound discharge with no other complications to a life-threatening condition. Other clinical outcomes of SSIs include poor scars that are cosmetically unacceptable, persistent pain and itching, restriction of movement and a significant impact on emotional wellbeing.

This report includes data captured both during the initial surgical procedure, revision surgery, and any other follow-up visits to the hospital (whether they were planned or not). The surveillance incorporates data collected by clinical teams and uses internationally agreed definitions [5], allowing Welsh data to be compared with and incorporated into other international databases, such as the ECDC European SSI database. This report details results obtained for surveillance data capture in 2018.

Data interpretation

Surgical site infection (SSI) rates in this report are calculated as the number of infections as a proportion of total procedures entered onto the web form. This is reported as a rate per 100 procedures.

 $SSI \ rate \ = \ \frac{number \ of \ SSI}{number \ of \ reported \ procedures} \times 100\%$

SSIs in orthopaedic procedures are most likely to occur within the first 90 days postprocedure. They can be reported later than 90 days and the system allows for surveillance of infections for up to five years following the procedure, however, an infection occurring more than a year following the procedure is rare, but is possible if the infection can be definitively linked to the surgery.

Extended surveillance only applies to electronically reported procedures, and older procedures reported on paper forms are generally monitored up to the post-discharge review at the surgeon's clinic.

All confidence intervals are set to 95% confidence unless stated otherwise.

Summary

NOTE: The information in this annual report may differ from that found in the more regular reports due to delays in data arriving at Public Health Wales and amendments being made following the issue of reports.



Streptococcus spp. are most frequently identified family of organisms Identified in <mark>83%</mark> of reported SSIs in 2018

SSI rate of 0.18% which is 9 times higher than the next most identified family of organisms

5,501 procedures were reported in 2018. Of these procedures, 12 were reported to have developed an SSI which equates to an SSI rate of 0.22%. Knee procedures had a higher SSI rate (0.37%) than hip procedures (0.12%).

Section 1: SSI rates

Annual SSI rates



Figure 1 – Graph showing the change in rate of SSI by year of procedure for primary hip procedures.



Figure 2 – Graph showing the change in rate of SSI by year of procedure for primary knee procedures.

The SSI rate has been gradually decreasing since 2008, however, there was a sharp decrease in primary knee procedures in 2016 when electronic data capture was introduced. This drop in rates is present for both hip and knee procedures, as well as the overall rates but most apparent in knee procedures.

Section 2: Frequency of revisions

Annual Revisions



Figure 3 – Graph showing the change in frequency of revision procedures by year for all hip procedures.



Figure 4 – Graph showing the change in frequency of revision procedures by year for all knee procedures.

Section 2: Procedure details and demographics

Table 1 – Procedure numbers and SSI rate breakdown by procedure type and demographic details.

Wales 2018				Wales 2008-2018					
	Procedures	SSI S	SSI rate		Procedures	SSI	SSI rate		
Overall	5,501	12	0.2%	Overall	59,534	644	1.1%		
Procedure types	5			Procedure types					
Primary hip	2,532	3	0.1%	Primary hip	26,310	256	1.0%		
Revision hip	154	0	0.0%	Revision hip	1,131	12	1.1%		
Primary knee	2,402	9	0.4%	Primary knee	29,669	359	1.2%		
Partial knee*	290	0	0.0%	Partial knee*	979	1	0.1%		
Revision knee	100	0	0.0%	Revision knee	808	11	1.4%		
Other/unknown	23	0	0.0%	Other/unknown	637	5	0.8%		
Age groups				Age groups					
Under 65	1,725	6	0.3%	Under 65	19,316	230	1.2%		
65-74	2,071	2	0.1%	65-74	22,193	224	1.0%		
75-84	1,467	3	0.2%	75-84	15,101	162	1.1%		
85 and over	204	1	0.5%	85 and over	2,265	19	0.8%		
Unknown	34	0	0.0%	Unknown	659	9	1.4%		
Sex				Sex					
Male	2,294	5	0.2%	Male	25,158	319	1.3%		
Female	3,147	6	0.2%	Female	34,016	321	0.9%		
Unknown	60	1	1.7%	Unknown	360	4	1.1%		
Report source				Report source					
Electronic	5,501	12	0.2%	Electronic	19,666	60	0.3%		
Paper forms	0	0		Paper forms	39,868	584	1.5%		
*Unicondylar and	patellofemoral kne	e replacen	nents.			2.57%			
						1.93%			
				Wales N	1ean 🚪	1.29%			
				2008-2	018	0.64%			
						0.00%			

Analysing data across different demographics enables the observation of potential risk factors associated with the development of an SSI, in this case the SSI rate for primary knee procedures appears to be four times higher to that of primary hip procedures in 2018. This observation does not hold across years, with 2017 seeing the reverse, with a primary hip SSI rate more than twice that of primary knee procedures (0.46% and 0.18% respectively).

There is a loose association between age and the risk of SSI across annual data, with those patients aged under 65 being at a greater risk than patients aged 65 and over, this trend is not apparent when looking at 2018 data only (P=0.029, P=0.068 for 2018 data only). There is also a significant difference in SSI rates between sexes (P=0.007), and men have a higher risk of developing an SSI than women. This trend is also present in 2018, however with lower numbers and therefore, less significance



A comparison in SSI rates between procedures reported on paper and procedures reported electronically will be made in the last section (Data quality) but, as can be seen in the above table, the SSI rate in the electronic data is still substantially lower than paper-based data. Individual hospital and health board SSI rates can be found in the appendix.

Figure 5 – Graph to show the proportion of different procedure types for Orthopaedic procedures conducted each year, 2008-2018.

It is evident that since the introduction of electronic data capture in 2015 that the numbers of revision knee procedures have increased, this is in contrast to revision hip procedures, which have seen relative increases and decreases at all times since 2008.

Procedure types

Joint	OPCS	Name	Procedures	SSI	SSI rate	(95% CI)
Нір	W371	Primary total hip replacement using cement	871	2	0.23%	(0.03 - 0.83)
	W373	Revision of total hip replacement using cement	30	0	0.00%	(0.00 - 11.57)
	W381	Primary total hip replacement not using cement	1034	1	0.10%	(0.00 - 0.54)
	W383	Revision of total hip replacement not using cement	110	0	0.00%	(0.00 - 3.30)
	W391	Primary total hip replacement NEC	296	0	0.00%	(0.00 - 1.24)
	W393	Revision of total hip replacement NEC	14	0	0.00%	(0.00 - 23.16)
	W581	Primary hip resurfacing	0	0	0.00%	(0.00 - 0.00)
Knee	W401	Primary total knee replacement using cement	2251	8	0.36%	(0.15 - 0.70)
	W403	Revision of total knee replacement using cement	ment using cement 95 0 0.00% (C			
	W411	Primary total knee replacement not using cement	mary total knee replacement not using cement 145 1 0.69%			
	W413	Revision of total knee replacement not using cement	5	0	0.00%	(0.00 - 52.18)
	W421	421 Primary total knee replacement NEC		0	0.00%	(0.01 - 24.71)
	W423	Revision of total knee replacement NEC	0	0	0.00%	(0.00 - 60.24)
	W428	Other total replacement of knee joint	4	0	0.00%	(0.00 - 13.72)
	Z844	Patellofemoral knee replacement	25 0 0.00%		(0.00 - 1.46)	
	Z845	Unicondylar knee replacement - medial	250	0	0.00%	(0.00 - 21.80)
	Z846	Unicondylar knee replacement - lateral	15	0	0.00%	(0.00 - 21.80)
Unknown	W05	Prosthetic replacement of bone	1	0	0.00%	(0.00 - 0.98)

Table 2 – Breakdown of procedure numbers and SSI rates for individual procedure types.

*Procedures reported in 2018 only

Mean SSI Rate	0.58%
	0.39%
	0.19%
	0.10%
	0.00%





Figure 6 – Graph showing the relative usage of prophylactic antibiotics in 2017 vs 2018.

Table 3 – Prophylactic antibiotics by frequency of use, with their associated SSI rates for 2018.

	Procedures	SSI	SSI rate	(95% CI)
Ceftriaxone + Teicoplanin	86	0	0.00%	(0.00% - 0.61%)
Cefuroxime	1087	0	0.00%	(0.00% - 12.34%)
Flucloxacillin + Gentamicin	182	0	0.00%	(0.11% - 1.49%)
Gentamicin	177	0	0.00%	(0.00% - 0.31%)
Gentamicin + Teicoplanin	3322	11	0.33%	(0.00% - 1.89%)
Teicoplanin	27	0	0.00%	(0.00% - 2.06%)
Other	32	1	3.13%	(0.17% - 0.59%)
Unknown	588	0	0.00%	(0.07% - 14.16%)

	0.58%
	0.39%
Mean SSI Rate	0.19%
	0.10%
	0.00%

Compared to 2017 the usage profile of prophylactic antibiotics has shifted, the use of Ceftriaxone and Teicoplanin has reduced and the combination of Gentamicin and Teicoplanin has increased as a result to become the most frequently used antibiotic combination for pre-surgery prophylaxis.

Section 3: Post-procedure details

Microbiology results

Table 4 – Rates of SSI by individual microorganism.

	2018		2014*-20	18
	Number of SSI	SSI rate	Number of SSI	SSI rate
Enterococcus	1	0.02%	13	0.05%
E. faecalis	0	0.00%	5	0.02%
E. faecium	1	0.02%	8	0.03%
Escherichia	1	0.02%	2	0.01%
E. coli	1	0.02%	2	0.01%
Klebsiella	1	0.02%	3	0.01%
K. pneumoniae	1	0.02%	2	0.01%
Pseudomonas	1	0.02%	1	0.00%
P. aureginosa	1	0.02%	1	0.00%
Staphylococcus	10	0.18%	43	0.16%
S. aureus	6	0.11%	28	0.11%
-No resistance data	6	0.11%	28	0.11%
S. (not specified)	2	0.04%	2	0.01%
Coagulase negative	2	0.04%	13	0.05%
-S. epidermidis	1	0.02%	7	0.03%
-S. haemolyticus	0	0.00%	5	0.02%
-Not specified	1	0.02%	1	0.00%

Other	1	0.02%	1	0.00%
Other Gram negative bacilli	1	0.02%	1	0.00%
No microbiology results	1	0.02%	699	2.63%

SSI rate shown as a percentage of procedures where the organism was present.

Tests can be positive for multiple organisms, therefore, some infections may be present multiple times under different organisms.

*Microbiology data was not collected prior to 2014

0.20%
0.19%
0.16%
0.08%
0.00%

Overall, the most common organism present in SSIs is *Staphylococcus sp.* Within this group, *S. aureus* is the most common single species, and this was present six times in 2018, which represents 50% of all infections in the year. It should be noted that while the frequency of *S. aureus* positive tests decreased, its relative proportion of infections has increased from 2017 at 10 times and 43% respectively. Resistance data was not provided for these cases, and it is therefore not possible to confirm whether these were methicillin sensitive or resistant. Coagulase negative Staphylococci as well as Enterococci were also relatively common.

Since the surveillance system captures multiple organisms, these numbers are higher than the total number of SSI, as some infections have multiple organisms recorded. There is also no way to indicate causation and, therefore, we are unable to confirm whether any of these organisms are the causative agent or are simply commensal organisms picked up along with the causative agent. There is a high probability that some Staphylococci and Enterococci in the surveillance system are purely commensal, and that others are the causative agent.

Section 4: Data quality

There are many inconsistencies in data reporting for the orthopaedic SSI surveillance scheme, with some hospitals reporting more consistently than others.

Most hospitals across Wales remain compliant in reporting Orthopaedic procedures that we have under surveillance, but a major issues arises in the fact that many of these hospitals have not reported any SSIs for periods ranging from a few months to several years. A lot of this disparity can be attributed to the 2015 switch over to electronic data reporting, where the change in procedures has meant that some hospitals are reporting most procedures on the system but are not updating these records when an infection is identified.

This issue was highlighted in 2017 and there have been steps taken this year to attempt to engage hospitals with reporting infections and to identify the issues that need to be dealt with in order to improve compliance. Morriston hospital has seen the largest relative gains in compliance going into the New Year, 2019.



Figure 7 – Proportions of data reporting sources per year, depicting the change from Paper to Electronic reporting sources since 2014.

Paper surveillance of orthopaedic procedures was discontinued in 2016 and, despite the switch, there has been no substantive change in the number of procedures being reported to PHW. There was a slight decrease in the number of procedures reported in 2018 compared to the previous year.

Discussion

In 2018, there were 5,501 procedures reported to Public Health Wales. Of these procedures, 12 were reported to have developed an SSI following the procedure, which equates to a Wales SSI rate of 0.22%.

It is difficult to ascertain the major risk factors contributing to the development of SSIs, however it appears as though there is a negative correlation between SSI rate and age with the under 65s having an SSI rate of 0.35%, and 50% of all SSIs reported this year being within this age group. The trend is not linear however, as a small numbers of procedures and a single SSI have inflated the SSI rate of the over 85s to 0.49%.

The gender disparity in SSI rate was not significant in 2018 with the female SSI rate at 0.22% compared to the male SSI rate at 0.19%. The 2018 data is a reversal of the 10 year trend which shows a significant slant towards males having a higher prevalence of SSIs at 1.27% compared to 0.94% for females.

In general, there is a lot of variability in reporting practices at hospitals. Some are fully compliant with the surveillance scheme while others are not reporting infections, and some not even reporting procedures. This means that comparing a hospital's infection rates to the Wales average is not recommended, as simply the act of reporting infections is enough for them to be above the average SSI rate for Wales. It is unlikely that the zero rates at some hospitals is a true reflection of the SSI rate, especially given the procedure numbers involved.

When discussing the volatility in SSI it is important to consider the effect of using electronic data capture. As can be seen from the SSI rates after 2014, the switch to electronic data capture may have contributed to a reduction in SSI rates as a result of under-reporting. It is also interesting to note the rise in reported revision procedures in the same period. Our data capture method does not list the reporting of revision procedures as an optional component in the way that it was presented when paper based reporting was the norm, this is likely to be the source of the observed increase.

The orthopaedic reports have been kept at a monthly reporting frequency. This has enabled several hospitals to review their reported procedures from a provided line listing for the following month in a timely manner, which has aided the quality of the orthopaedic dataset. Unfortunately, this line listing does not seem to have acted as the necessary prompt required to trigger infection reporting generally.

The HARP team are committed to the improvement of the Orthopaedic surveillance system and therefore have been investigating a variety of methods to capture infection data from hospitals in a heuristic system that will limit the work hours required to produce an SSI rate for each hospital included. This system will act as an interim solution until a more permanent system is developed. While we go through the process of developing and implementing these new methods, we would appreciate it if hospitals could continue reporting via the usual channels.

References

- 1. Plowman R. The socioeconomic burden of hospital acquired infection. Euro surveillance: bulletin Européen sur les maladies transmissibles. 2000; 5(4).
- 2. Jenks PJ, Laurent M, McQuarry S, Watkins R. Clinical and economic burden of surgical site infection (SSI) and predicted financial consequences of elimination of SSI from an English hospital. Journal of Hospital Infection. 2014; 86(1): p. 24-33.
- 3. National Institute for Health and Care Excellence. Surgical site infections: prevention and treatment. [Online].; 2017 [cited 2018 June 1. Available from: https://www.nice.org.uk/guidance/CG74/chapter/1-Guidance.
- 4. Sykes PK, Brodribb RK, McLaws ML, McGregor A. When continuous surgical site infection surveillance is interrupted: The Royal Hobart Hospital experience. American Journal of Infection Control. 2005; 60(2): p. 422-427.
- 5. Horan TC, Gaynes RP, Martone WJ, Jarvis WR, Emori TG. CDC definitions of nosocomial surgical site infections, 1992: A modification of CDC definitions of surgical wound infections. Infection Control & Hospital Epidemiology. 1992; 13(10): p. 606-608.
- 6. van Walraven C, Musselman R. The Surgical Site Infection Risk Score (SSIRS): A Model to Predict the Risk of Surgical Site Infections. PLoS One. 2013; 8(6): p. e67167.

Appendix

	Hip Procedures		Knee Procedures			All Procedures			
	Procedures	SSI	SSI rate*	Procedures	SSI	SSI rate*	Procedures	SSI	SSI rate*
Abertawe Bro Morgannwg UHB	83	0		105	0		188	0	
Aneurin Bevan UHB	841	0		1083	6	0.55%	1926	6	0.31%
Betsi Cadwaladr UHB	695	0		590	0		1285	0	
Cardiff and Vale UHB	46	0		43	0		89	0	
Cwm Taf UHB	373	2	0.54%	413	3	0.73%	787	5	0.64%
Hywel Dda UHB	656	1	0.15%	569	0		1226	1	0.08%

Table 5 – Health board SSI rates in 2018, broken down by joint.

	0.78%
	0.39%
Mean SSI rate	0.19%
	0.10%
	0.00%

Table 6 – Health board level microbiology data 2018

	ABMUHB		ABUHB		BCUHB		CVUHB		СТИНВ		HDUHB	
	Number of SSI	SSI rate										
Enterococcus	0	0.00%	1	0.02%	0	0.00%	0	0.00%	0	0.00%	0	0.00%
E. faecalis	0	0.00%	0	0.00%	0	0.00%	0	0.00%	0	0.00%	0	0.00%
E. faecium	0	0.00%	1	0.02%	0	0.00%	0	0.00%	0	0.00%	0	0.00%
Escherichia	0	0.00%	1	0.02%	0	0.00%	0	0.00%	0	0.00%	0	0.00%
E. coli	0	0.00%	1	0.02%	0	0.00%	0	0.00%	0	0.00%	0	0.00%
Klebsiella	0	0.00%	1	0.02%	0	0.00%	0	0.00%	0	0.00%	0	0.00%
K. pneumoniae	0	0.00%	1	0.02%	0	0.00%	0	0.00%	0	0.00%	0	0.00%
Pseudomonas	0	0.00%	0	0.00%	0	0.00%	0	0.00%	1	0.02%	0	0.00%
P. aureginosa	0	0.00%	0	0.00%	0	0.00%	0	0.00%	1	0.02%	0	0.00%
Staphylococcus	0	0.00%	3	0.05%	0	0.00%	0	0.00%	6	0.11%	1	0.02%
S. aureus	0	0.00%	2	0.04%	0	0.00%	0	0.00%	4	0.07%	0	0.00%
-No resistance data	0	0.00%	2	0.04%	0	0.00%	0	0.00%	4	0.07%	0	0.00%
S. (not specified)	0	0.00%	0	0.00%	0	0.00%	0	0.00%	1	0.02%	1	0.02%
Coagulase negative	0	0.00%	1	0.02%	0	0.00%	0	0.00%	1	0.02%	0	0.00%
-S. epidermidis	0	0.00%	1	0.02%	0	0.00%	0	0.00%	0	0.00%	0	0.00%
-S. haemolyticus	0	0.00%	0	0.00%	0	0.00%	0	0.00%	0	0.00%	0	0.00%
-Not specified	0	0.00%	0	0.00%	0	0.00%	0	0.00%	1	0.02%	0	0.00%
						_				-		
Other	0	0.00%	0	0.00%	0	0.00%	0	0.00%	1	0.02%	0	0.00%
Other Gram negative bacilli	0	0.00%	0	0.00%	0	0.00%	0	0.00%	1	0.02%	0	0.00%
No microbiology results	0	0.00%	1	0.02%	0	0.00%	0	0.00%	0	0.00%	0	0.00%

SSI rate shown as a percentage of procedures where the organism was present.

Tests can be positive for multiple organisms, therefore, some infections may be present multiple times

under different organisms.

*Microbiology data was not collected prior to 2014

0.029% 0.022% 0.015% 0.007% 0.000%