

Rhaglen Sgrinio
Ymlediaid Aortig
Abdomenol Cymru



Wales Abdominal
Aortic Aneurysm
Screening Programme




GIG
CYMRU
NHS
WALES

Iechyd Cyhoeddus
Cymru
Public Health
Wales

Wales Abdominal Aortic Aneurysm Screening Programme

Annual Statistical Report 2019-20

October 2021



Screening Division of Public Health Wales Publication

About us

Public Health Wales exists to protect and improve health and wellbeing and reduce health inequalities for people in Wales.

We are part of the NHS and report to the Minister for Health and Social Services in the Welsh Government.

Our vision is for a healthier, happier and fairer Wales. We work locally, nationally and, with partners, across communities in the following areas:

Health protection – providing information and advice and taking action to protect people from communicable disease and environmental hazards

Primary, community and integrated care – strengthening its public health impact through policy, commissioning, planning and service delivery

Microbiology – providing a network of microbiology services which support the diagnosis and management of infectious diseases

Safeguarding - providing expertise and strategic advice to help safeguard children and vulnerable adults

Screening – providing screening programmes which assist the early detection, prevention and treatment of disease

Health intelligence – providing public health data analysis, evidence finding and knowledge management

NHS quality improvement and patient safety – providing the NHS with information, advice and support to improve patient outcomes

Policy, research and international development – influencing policy, supporting research and contributing to international health development

Health improvement – working across agencies and providing population services to improve health and reduce health inequalities

Further information

Web: phw.nhs.wales
Email: general.enquiries@wales.nhs.uk
Twitter: [@PublicHealthW](https://twitter.com/PublicHealthW)
Facebook: facebook.com/PublicHealthWales

This report is a detailed summary of information on work undertaken by the Wales Abdominal Aortic Aneurysm Screening Programme for the year from April 2019 to the end of March 2020.

Publication Details:

Title: Wales Abdominal Aortic Aneurysm Screening Programme Annual Statistical Report 2019-20

Date: This report published October 2021

ISBN: 978-1-78986-154-400

For more information about this report contact:

Helen Clayton, Lead Informatics and Data Services Manager,
Informatics Division, Floor 6, Public Health Wales, Number 2 Capital Quarter,
Tyndall Street, Cardiff, CF10 4BZ
Tel: 029 2010 4405

Email: screening.information@wales.nhs.uk

Rydym yn croesawu gohebiaeth a galwadau ffôn yn Gymraeg. Byddwn yn ateb gohebiaeth yn Gymraeg heb oedi / We welcome correspondence and phone calls in Welsh. We will respond to correspondence in Welsh without delay.

Quality Assurance Statement

Screening data records are constantly updated. The databases used by Public Health Wales Screening Division are updated on a daily basis when records are added, changed or removed (archived). This might relate to when a person has been identified as needing screening; has had screening results that need to be recorded, or has a change of status and no longer needs screening respectively. Data is received from a large number of different sources with varying levels of accuracy and completeness. The Screening Division checks data for accuracy by comparing datasets – for example GP practice data – and corrects the coding data where possible. It should be noted that there are sometimes delays in data collection – for example a person might not immediately register with their GP if they move address. These delays will therefore affect the completeness of the data depending on individual circumstances. In addition, the reader should be aware that data is constantly updated and there might be slight readjustments in the numbers cited in this document year on year because of data refreshing. We occasionally suppress numbers lower than five when the data is potentially sensitive.

© 2021 Public Health Wales NHS Trust

Material contained in this document may be reproduced under the terms of the Open Government Licence (OGL)

www.nationalarchives.gov.uk/doc/open-government-licence/version/3/ provided it is done so accurately and is not used in a misleading context.

Acknowledgement to Public Health Wales NHS Trust to be stated. Copyright in the typographical arrangement, design and layout belongs to Public Health Wales NHS Trust.

This document is also available in Welsh.

Contents

1	INTRODUCTION	6
1.1	'KEY MESSAGES' FOR THE PUBLIC.....	6
1.2	Programme delivery	8
1.3	Screening pathway.....	8
2	HEADLINE STATISTICS	9
3	DATA.....	10
3.1	Uptake	10
3.2	Non-visualised.....	15
3.3	Men who self-refer	16
3.4	Abdominal aortic aneurysms detected	17
3.5	AAA surveillance uptake.....	18
3.6	Referral to multi-disciplinary team	18
4	DEFINITIONS.....	19
5	PRODUCTION TEAM	20

Tables and Graphs

Table 1a:	AAA screening uptake by health board of residence	10
Table 1aa:	AAA screening uptake by month of invite.....	11
Graph 1a:	AAA screening uptake by health board of residence (%).....	12
Table 1b:	AAA screening uptake by deprivation quintile and health board of residence (%).....	13
Graph 1b:	AAA screening uptake by deprivation quintile and health board of residence (%)	14
Table 2:	Non-visualised rate by health board of residence	15
Table 3:	Number of those screened that have an AAA (≥ 3 cm) detected by health board of residence	17

1 Introduction

This is the sixth annual statistical report published by the Wales Abdominal Aortic Aneurysm Screening Programme (WAAASP).

WAAASP was launched in May 2013 and aims to halve the number of abdominal aortic aneurysm (AAA) related mortality by 2025 in the eligible population through a systematic screening programme for 65 year old men resident in Wales. Since 1 May 2015, men who have never been for an NHS AAA screening scan and who have not been offered AAA screening as they turned 65 before it was available in Wales can contact the local screening offices to request an AAA screening scan.

Research evidence has shown that a high quality screening programme for AAA can reduce deaths from ruptured aortic aneurysm by around 50% in men aged 65 – 74 years¹. In February 2007, the UK National Screening Committee approved the introduction of AAA screening for men aged 65 using abdominal ultrasound scanning provided:

- Invited men were given clear information about the risks of elective surgery, and
- Vascular networks were in place to treat individuals referred from screening

In March 2020, the difficult decision was taken to pause some of the national screening programmes in Wales, including WAAASP, in response to the COVID-19 pandemic. The decision to pause primary screening affected clinics scheduled for after March 17th, and the decision to pause self referrals was taken on March 24th. Although this only affected a very small proportion of the year 2019-20, the pause will have affected some measures such as uptake. We are also aware that the emerging pandemic from January – March did affect the behaviour of some of those invited for screening with some men opting to defer or cancel their appointment.

1.1 'Key messages' for the public

- Undertaking the abdominal aortic aneurysm (AAA) screening test reduces the risk of dying from an AAA. Finding an AAA early gives the man the best chance of treatment and survival

¹ Ashton HA, Buxton MJ, Day NE, Kim LG, Marteau TM, Scott RAP et al. (2002) Multicentre Aneurysm Screening Study Group. The Multicentre Aneurysm Screening Study (MASS) into the effect of abdominal aortic aneurysm screening on mortality in men: a randomised controlled trial. *Lancet*;360 (9345):1531-9

- The aorta is the main blood vessel that supplies blood to the body. An AAA is a swelling of the aorta in the abdomen, which left undetected, may split or rupture
- AAA screening involves a simple ultrasound scan to measure the abdominal aorta
- AAA screening is a free NHS test carried out in community clinics
- Taking part in AAA screening is the man's choice

1.2 Programme delivery

The Screening Division of Public Health Wales is responsible for managing, delivering and quality assuring the programme. The programme employs a Head of Programme, Programme Manager, Quality Assurance Vascular Surgical Lead, Clinical Imaging Advisor, Quality, Education and Training Lead, three clinical skills trainers and an All-Wales Administration Coordinator with support from a secretarial and administration team. Although an all-Wales programme, there is regional coordination by three Regional Coordinators and a team of 19 screeners who deliver the screening in the community.

1.3 Screening pathway

- 65 year old men resident in Wales and registered with a GP are invited for a one-off ultrasound scan to check whether they have an AAA
- The test involves a simple scan of the abdominal aorta, measuring the widest part of the aorta
- Ultrasound scanning is performed in 66 community clinics throughout Wales, including community hospitals, health clinics, primary resource centres and GP practices. Screening is undertaken in HMP Berwyn, HMP Parc, HMP Usk and HMP Prescoed
- Men with an abdominal aortic diameter of less than 3cm are discharged from the programme
- Men with a small or medium AAA are included in the surveillance programme and are offered:
 - small AAA (3 - 4.4cm) an annual scan
 - medium AAA (4.5 - 5.4cm) a scan every three months
 - a phone appointment with the AAA surveillance nurse to discuss the result and its health implications
 - encouragement to make an appointment with their GP for lifestyle and health advice, blood pressure monitoring and best medical therapy
- Men with a large AAA of 5.5cm or more (or a growth of 1cm or more in 12 months) are referred to the regional elective Vascular Network Multi-professional team (Multi-disciplinary Team or MDT)
- Men with a non-visualised aorta are usually offered a second appointment with WAAASP. If the second appointment is unsuccessful, the man is referred to a medical imaging department to measure his abdominal aorta

More information is available at www.aaascreening.wales.nhs.uk

2 **Headline statistics**

This report covers the time period from April 2019 to March 2020. Due to the Covid-19 pandemic AAA screening was paused on 18th March 2020.

Uptake is defined as those invited in the year 2019-20 receiving a scan by 30 June 2020. Uptake has been significantly affected by the pause in AAA screening. As no clinics were operating from March 18th, men invited towards the end of the financial year who did not attend their first appointment, had reduced opportunity to reschedule or be invited for a second appointment.

- The uptake for participants invited between April 2019 and March 2020 was 71.9%, ranging from 63.4% in Powys Teaching Health Board to 76.7% in Betsi Cadwalader University Health Board
- Uptake figures are higher in those men living in the least deprived areas (77.4%) compared to the most deprived areas (64.7%)
- Having increased in 2018-19, uptake of AAA screening decreased in 2019-20 to below the 80% target at 71.9% although this reflects the impact of the COVID pandemic on the last few months of the financial year.

In April 2019 – March 2020:

- 17,045 eligible men were invited by the programme from April 2019 to March 2020. Of these, 12,256 men attended for their first WAAASP scan and had a definitive scan result
- Of the men who attended for their screening from April 2019 to March 2020, 120 men had an AAA (0.9%) detected by the screening programme
- 53 men scanned needed a referral to the elective vascular network MDT. 88.7% of men were referred within two working days of the scan being taken
- 50 men had open or endovascular surgery. This is a different cohort to the men who were scanned and referred in the year. 12 (24.0%) of these had their surgery completed within four or eight weeks of the referral being received, depending on the size of the AAA detected

- 1121 (91.2%) surveillance scans were taken within standard from a possible 1229 opportunities (medium AAA on quarterly surveillance within 11 to 15 weeks, small AAA on annual surveillance within 50 to 56 weeks of their previous successful scan)
- 922 self-referred men were screened with 20 AAA (2.2%) detected

3 Data

3.1 Uptake

Standard: A minimum of 80% of invited men attending AAA screening are tested.

At an all-Wales level, uptake for 2019/20 was 71.9%. This is a significant decrease from 80.8% in 2018/19 when the standard was met. This decrease reflects the impact of the COVID pandemic on uptake, particularly in March of 2020 when the programme paused during the month and clinics were cancelled, but also for men invited in January and February of that year too.

Table 1a: AAA screening uptake by health board of residence

Health Board	Invited	Tested	% Uptake
Aneurin Bevan University	3308	2397	72.5
Betsi Cadwalader University	3893	2986	76.7
Cardiff & Vale University	2256	1557	69.0
Cwm Taf Morgannwg University	2404	1728	71.9
Hywel Dda University	2236	1572	70.3
Powys Teaching	922	585	63.4
Swansea Bay University	1731	1209	69.8
Unknown	295	222	75.3
All Wales	17045	12256	71.9

Note: uptake stated (of those eligible and invited in the year, number tested by 30 June 2020). Unknown refers to men who cannot be allocated to a health board however they are included in the all-Wales total.

The following table shows uptake each month throughout the year. It demonstrates consistent good performance at the start of the year, followed by low uptake at the start of 2020.

Table 1aa: AAA screening uptake by month of invite

Year	Month	Invited	Tested	% Uptake
2019	April	1506	1229	81.6
2019	May	1759	1424	81.0
2019	June	1666	1354	81.3
2019	July	1725	1371	79.5
2019	August	1222	975	79.8
2019	September	1412	1137	80.5
2019	October	1506	1201	79.7
2019	November	1527	1182	77.4
2019	December	929	691	74.4
2020	January	1628	1059	65.0
2020	February	1463	626	42.8
2020	March	702	7	1.0
Total		17045	12256	71.9



Graph 1a: AAA screening uptake by health board of residence (%)

Table 1b: AAA screening uptake by deprivation quintile and health board of residence (%)

Health Board	Q1 – least deprived	Q2	Q3	Q4	Q5 – most deprived	Total
Aneurin Bevan University	78.6	76.1	74.1	71.4	63.0	72.5
Betsi Cadwalader University	79.2	77.6	76.9	73.4	74.2	76.7
Cardiff & Vale University	77.3	68.1	63.7	65.8	57.4	69.0
Cwm Taf Morgannwg University	79.2	77.0	70.2	72.7	65.8	71.9
Hywel Dda University	73.5	68.0	72.0	70.5	66.2	70.3
Powys Teaching	76.8	61.2	66.3	56.1	55.6	63.4
Swansea Bay University	72.4	70.3	69.7	72.6	64.6	69.8
Unknown	-	-	-	-	-	75.3
All Wales	77.4	71.9	72.3	71.4	64.7	71.9

Note: Unknown refers to men who cannot be allocated to a health board however they are included in the all-Wales total.

Graph 1b: AAA screening uptake by deprivation quintile and health board of residence (%)



This shows that, in general across all the health boards, uptake decreases as deprivation score increases. It should be noted that Quintile 5 in Powys is composed of small numbers.

3.2 Non-visualised

Standard: $\leq 3\%$ of consented appointments resulting in a non-visualised aorta.

Table 2: Non-visualised rate by health board of residence

Health Board	Scans	Non-visualised	Non-visualised Rate (%)
Aneurin Bevan University	3,051	38	1.2
Betsi Cadwalader University	3,869	31	0.8
Cardiff & Vale University	1,983	28	1.4
Cwm Taf Morgannwg University	2,075	50	2.4
Hywel Dda University	2,076	15	0.7
Powys Teaching	7,64	9	1.2
Swansea Bay University	1,593	10	0.6
Unknown	289	5	1.7
All Wales	15,700	186	1.2

Note: non-visualised data refers to the number of completed appointments where the abdominal aorta was not seen.

3.3 Men who self-refer

Since 1 May 2015, men over 65 who have not received an NHS ultrasound screening scan for AAA can self-refer by contacting the screening programme to request an appointment. It is anticipated that the number of men self-referring for AAA screening will decline as the programme matures.

During 2019-20, there were 922 self-referred men scanned with 20 (2.2%) AAA detected. This only includes men who have not previously been invited by the programme.

3.4 Abdominal aortic aneurysms detected

Standard: Of those screened, it is expected that 1% will have an AAA ($\geq 3\text{cm}$).

Table 3: Number of those screened that have an AAA ($\geq 3\text{cm}$) detected by health board of residence

Health Board	Attended	AAA Total	Detection Rate (%)
Aneurin Bevan University	2,632	20	0.8
Betsi Cadwalader University	3,255	37	1.1
Cardiff & Vale University	1,701	14	0.8
Cwm Taf Morgannwg University	1,855	11	0.6
Hywel Dda University	1,747	23	1.3
Powys Teaching	654	2	0.3
Swansea Bay University	1,323	11	0.8
Unknown	238	2	0.8
All Wales	13,405	120	0.9

Note: Men with AAA ($\geq 3\text{cm}$) detected only counted on first definitive scan not surveillance scans. Non-visualised is not a definitive scan result. Unknown refers to men who cannot be allocated to a health board however; they are included in the all-Wales total.

3.5 AAA surveillance uptake

The surveillance uptake for this time period 2019-20, includes both men with a medium AAA detected, who are invited for quarterly surveillance, and men with a small AAA detected, who are invited for annual surveillance.

During 2019-20, 1,121 surveillance appointments were attended (91.2%) within standard from a possible 1,229 opportunities (men with medium AAA on quarterly surveillance should be re-scanned within 11 to 15 weeks, and men with small AAA on annual surveillance should be re-scanned within 50 to 56 weeks of their previous successful scan).

3.6 Referral to multi-disciplinary team

During 2019-20, 53 men were scanned and needed a referral to the elective vascular network MDT. This does not include referrals to on call vascular services (i.e. those with a very large AAA detected). Of the total referred, 88.7% were referred within two working days of the scan being taken.

50 men had open or endovascular surgery. This is a different cohort to the men who were scanned and referred in the year. 12 (24.0%) of these had their surgery completed within four or eight weeks of the referral being received, depending on size of AAA detected. Compliance with this timeliness standard has been discussed at the joint WAAASP and EVN MDT Coordinators meetings. The MDT coordinators submit an exception report for all men who breach the timeliness of repair standard and/ or have the repair in a spoke hospital rather than the agreed centralisation site.

There is a slight decrease in compliance of this standard from the previous Annual Statistical Report. In 2018-19, 12 men (24.5%) had their surgery within the timeliness standard. The reasons for the delay in treatment during both years are multifactorial and include:

- Men with co-morbidities
- Reduction in theatre capacity
- Delays in pre-operative diagnostic tests
- Variation in progress in the development of the regional elective vascular networks

Covid-19 pandemic; during the first wave of the pandemic the Vascular Society for Great Britain and Ireland recommended that elective AAA repair should only be undertaken if AAA was larger than 7cm. This affected the

timeliness of AAA repair for men referred during the latter months of 2019-20.

4 Definitions

This section provides further detail on the calculations used in this report.

Eligible

For uptake calculations, eligible men were those resident in Wales and were invited in the time period. Men who were registered manually (such as self-referrals) are excluded. Men invited who were ceased from the programme in the time period due to being out of cohort are removed.

Uptake

Men were counted as having responded to their invitation if they were invited during the April – March time period and attended by 30 June 2020.

Deprivation

Deprivation quintiles were assigned using the Welsh Index of Multiple Deprivation (WIMD) 2014, measured at lower super output area (LSOA) level. LSOAs are ranked into quintiles at an all-Wales level so they can be compared between health boards. This means that there will not be an equal proportion of people in each quintile when you look at each health board e.g. in Monmouthshire, 40% of the population live in the least deprived quintile of Wales but no areas fall into the Welsh most deprived quintile.

Health board

This is health board of residence.

Result

A definitive scan result excludes those where the final outcome is that the abdominal aorta could not be visualised.

5 Production team

The production team for this report are all employed within Public Health Wales and are listed below.

Jeremy Surcombe	Head of Wales Abdominal Aortic Aneurysm Screening Programme
Dr Sharon Hillier	Director of Screening Division
Sikha de Souza	Consultant in Public Health
Helen Clayton	Lead Informatics and Data Services Manager
Richard Wakely	Senior Informatics and Data Analyst
Guy Stevens	Deputy Informatics and Data Services Manager
Rebecca Turner	Informatics and Data Analyst
Hannah Lindsay	Communications Manager
Diane Rawlings	Personal Assistant
Rhys George	Cofus CTF (Translator)