

This information is for you if your fetal anomaly scan has shown that your baby has renal pelvis dilatation (RPD)

The information applies if RPD is the only unexpected finding on the scan.


Every pregnant woman in Wales is offered a fetal anomaly scan when they are between 18 and 20 weeks pregnant. The scan is a way of checking that your baby appears to be developing as expected. This means it may show up unexpected findings that would then need to be checked by further scans.

Renal pelvis dilatation

Renal pelvis dilatation is also known as RPD. RPD is found in less than 1 in 100 (less than 1%) anomaly scans performed in Wales.

What is RPD?

The renal pelvis is the area in your baby's kidney where urine collects. If the renal pelvis looks wider (more dilated) than usual, the sonographer will measure it. If it measures over 5 millimetres (mm), this is classed as dilated. RPD can be seen in just one kidney (unilateral) or both kidneys (bilateral).



What will happen next?

You will be seen by a midwife who specialises in antenatal screening or your hospital doctor (obstetrician) who will explain what care you will be offered next.

What care will I be offered?

If the RPD measured between 5.1mm and 10.0mm, you will be offered a further scan when you are 30 to 32 weeks pregnant.

If the measurement was 10.1mm or above, this is known as hydronephrosis and you will be given an appointment with an obstetrician who will discuss this in more detail with you. You will also be offered a further scan when you are 30 to 32 weeks pregnant.

What will the next scan tell me?

If the next measurement is 7.0mm or below

If the measurement following the next scan is 7.0mm or below, there will be no need for any further follow-up. In Wales this happens in around 8 out of 10 (80%) babies. These babies have about the same chance of going to hospital with a kidney problem during childhood as babies who did not have RPD seen on the anomaly scan.

If the next measurement is 7.1mm or above

If the scan at 30 to 32 weeks shows that the RPD measures 7.1mm or above, an appointment will be made for you to discuss this with a doctor who will explain this in more detail.

In Wales, in around 2 out of 10 (20%) babies with RPD at the anomaly scan, the renal pelvis remains dilated. For these babies, the risk of going to hospital with a kidney problem during childhood is around 20 times more likely than for babies with no RPD on the anomaly scan (38 in 100 children compared with 2 in 100 children).

The doctor will arrange for your baby to be seen by a paediatrician (baby doctor) after your baby is born.

If my baby needs a follow-up after they are born, what does this mean?

The paediatrician may offer you antibiotics for your baby to prevent a possible infection. They will also arrange another scan for when your baby is around two weeks old to see if the renal pelvis remains wider (dilated).



Further information

You can get more information from the hospital midwife who specialises in antenatal screening or your hospital doctor (obstetrician).

Antenatal Results and Choices (ARC)

Website: www.arc-uk.org

Phone: 0845 0772290 or 0207 7137486

Information from CARIS (Congenital Anomaly Register and Information System) and The Welsh Study of Mothers and Babies has been used in this leaflet to describe how RPD may present in your baby.

© Copyright 2022 Public Health Wales NHS Trust. All rights reserved.
Not to be reproduced in whole or part without the permission of the
copyright owner.

phw.nhs.wales/antenatal-screening

July 2022

RPD 4th Edition

