



COVID-19 vaccination schedule

Individuals not at greater risk of serious disease from COVID-19 infection

Age (years)	0-4	5-11	12-15	16-17	18-39	40+
Primary course 	Vaccination not currently recommended		2 doses Comirnaty® (PfizerBioNTech) >12 week interval ⁴		2 doses Comirnaty® (PfizerBioNTech) or Moderna (Spikevax®) ² 8 week interval	2 doses Comirnaty® (PfizerBioNTech) Moderna (Spikevax®) ² AstraZeneca (Vaxzevria®) ⁵ 8 week interval
Booster 	✗	✗	✗ No Booster not recommended	✔ Yes Comirnaty® (PfizerBioNTech) At least 3 month interval	✔ Yes Comirnaty® (PfizerBioNTech) Moderna (Spikevax®) ² At least 3 month interval	
Infection 	✗	✗	Wait at least 12 weeks between COVID-19 infection and COVID-19 vaccination ⁴		Wait 4 weeks between COVID-19 infection and COVID-19 vaccination	

Individuals who are at greater risk of serious disease from COVID-19 infection³

Age (years)	0-4	5-11	12-15	16-17	18-39	40+
Primary course 	Vaccination not currently recommended	2 doses ¹ Comirnaty® [▼] (Children 5-11yrs) (PfizerBioNTech) ⁶ 8 week interval	2 doses ¹ Comirnaty® (PfizerBioNTech) 8 week interval		2 doses ¹ Comirnaty® (PfizerBioNTech) or Moderna (Spikevax®) ² 8 week interval	2 doses ¹ Comirnaty® (PfizerBioNTech) Moderna (Spikevax®) ² AstraZeneca (Vaxzevria®) 8 week interval
Booster 	✗	✗ No Booster not recommended		✔ Yes Comirnaty® (PfizerBioNTech) At least 3 month interval	✔ Yes Comirnaty® (PfizerBioNTech) Moderna (Spikevax®) ² At least 3 month interval	
Infection 	✗	Wait 4 weeks between COVID-19 infection and COVID-19 vaccination				

¹3rd dose is advised if the person was severely immunocompromised at the time of first or second dose (refer to green book for details)

² The priming dose of Moderna Spikevax® COVID-19 Vaccine is 100 micrograms (0.5mL), The booster dose is 50 micrograms (0.25mL)

³ Health care, social care or care home staff, clinical risk groups, and those who are severely immunosuppressed or their household contacts

⁴ This interval may be reduced to 8 weeks in periods of high incidence or where there was concern about vaccine effectiveness

⁵ Vaxzevria should only be used in exceptional circumstances

⁶ Children aged 5 to 11 should receive two 10µg doses of Comirnaty®[▼] (Children 5-11yrs) (PfizerBioNTech)

