

**Current level of influenza activity:** *Circulation at low levels.*

**Trend:** *Decreasing*

**Confirmed cases since 2018 week 40: 2,451** (99.5% influenza A and 0.5% influenza B. Of influenza A cases, 54.4% were A(H1N1)pdm09, 16.6% were A(H3) and 29.1% were untyped)

## Key points – Wales


### Surveillance indicators suggest that influenza is circulating in Wales.

The sentinel GP consultation rate for influenza-like illness (ILI) has decreased and remained below baseline levels during week 11 (ending 17/03/2019), although influenza continues to be confirmed in patients with ILI, particularly those attending hospitals. Influenza was the most commonly detected cause of Acute Respiratory Infection (ARI) in hospital and non-sentinel GP patients, with 83 confirmed cases. Influenza A(H3N2) was dominant during week 11, although A(H1N1)pdm09 remains dominant for the season overall. Other causes of ARI, including rhinovirus, coronavirus, human metapneumovirus, adenovirus, and parainfluenza also continue to be detected.

- The Sentinel GP consultation rate for influenza-like illness (ILI) in Wales during week 11 was 7.5 consultations per 100,000 practice population. The consultation rate was highest in patients aged 25-34 years (11.9 per 100,000 practice population) (Table 1).
- The ILI consultation rate decreased compared to week 10 (8.7 per 100,000), and remains below baseline levels (Figure 1). Based on sentinel ILI consultation data from 2010 to 2018, an average influenza season in Wales lasts 14 weeks. This season, the sentinel ILI consultation rate crossed the baseline threshold in 2018 week 52 and returned to baseline levels in 2019 week 08 (9 weeks in total), however influenza continues to be detected in samples from patients with ILI and ARI as at week 11.
- The total number of consultations with Out of Hours (OOH) doctors in Wales reported to Public Health Wales during week 11 was 10,279. The proportion of respiratory-related consultations with OOH doctors increased from 18.1% to 18.6% (Figure 5). The percentage of calls to NHS Direct Wales which were 'influenza-related' (cold/flu, cough, fever, headache and sore throat) during week 11 decreased to 15.3% (Figure 6).
- No surveillance samples from patients with ILI, collected by sentinel GPs during week 11, had been received by Public Health Wales Microbiology as at 20/03/2019 (Figure 3).
- During week 11, 344 specimens were received and tested by Public Health Wales Microbiology from hospitalised and non-sentinel GP patients with acute respiratory symptoms. The following numbers of patients tested positive: 53 influenza A(H3N2), 13 influenza A(H1N1)pdm09, 17 influenza A (not typed), 48 rhinovirus, 22 coronavirus, 20 human metapneumovirus, 16 adenovirus, 13 for parainfluenza, three for enterovirus, one for RSV and one for mycoplasma (Figure 4). The proportion of samples from hospital patients positive for influenza decreased to 24.1%.
- During week 11, there were three outbreaks of acute respiratory illnesses (ARI) reported to Public Health Wales Health Protection teams, two outbreaks in a hospital (one confirmed as influenza (not typed)), and one outbreak in a residential home (where both influenza A(H1N1)pdm09 and influenza A(H3N2) were detected).
- At the end of week 11, uptake of influenza vaccine was: 68.2% in those aged 65 year and older, 43.8% in patients aged six months to 64 years at clinical risk, and 49.3% in children aged two and three years. In the 1,373 primary schools visited so far as part of the universal childhood influenza programme, uptake was 69.8%.


## Influenza activity in Wales

**Figure 1. Clinical consultation rate per 100,000 practice population in Welsh sentinel practices (as of 20/03/2019).**



\* The Moving Epidemic Method has been adopted by the European Centre for Disease Prevention and Control to calculate thresholds for GP ILI consultations for seasonally expected influenza activity in a standardised approach across Europe. The threshold calculated for Wales ILI consultation rates is 10.8 per 100,000. MEM thresholds used in this chart are based on influenza from 2010-11 to 2017-18 seasons.

**Figure 2. Clinical consultation rate per 100,000 practice population in Welsh sentinel practices (week 47 1996 – week 11 2019).**




\* Reporting changed to Audit+ surveillance system

**Table 1. Age-specific consultations (per 100,000) for influenza in Welsh sentinel practices, week 06 – week 11 2019 (as of 20/03/2019).**


Age group	6	7	8	9	10	11
< 1	0.0	0.0	0.0	0.0	0.0	0.0
1 - 4	6.2	12.4	6.2	0.0	6.2	0.0
5 - 14	9.2	4.6	2.3	0.0	0.0	2.3
15 - 24	27.7	8.5	23.5	6.4	10.7	8.6
25 - 34	15.9	17.9	25.9	10.0	13.9	11.9
35 - 44	46.2	27.3	23.1	14.7	8.4	10.5
45 - 64	27.9	29.8	24.2	14.0	13.0	6.5
65 - 74	15.1	6.5	6.5	4.3	6.5	8.6
75+	2.5	10.1	10.1	2.5	2.5	7.6
<b>Total</b>	<b>21.4</b>	<b>17.2</b>	<b>17.5</b>	<b>8.2</b>	<b>8.7</b>	<b>7.5</b>

**Figure 3. Specimens submitted for virological testing by sentinel GPs as of 20/03/2019, by week of sample collection, week 11 2018 - week 11 2019.**



\* Tested negative for influenza, adenovirus, rhinovirus, RSV, parainfluenza, mycoplasma, human metapneumovirus, enterovirus, bocavirus and coronaviruses.


**Figure 4. Specimens submitted for virological testing for hospital patients and non-sentinel GPs as of 20/03/2019 by week of sample collection, week 11 2018 – week 11 2019.**




Combined data for tests carried out in Public Health Wales Microbiology: Cardiff laboratory, provided by Public Health Wales Microbiology Cardiff Specialist Virology Centre.

### Out of Hours consultations and calls to NHS Direct Wales

**Figure 5. Weekly total consultations to Out of Hours services in Wales and numbers of respiratory-related diagnoses (as of 20/03/2019).**



**Figure 6. Influenza related calls to NHS Direct Wales<sup>1</sup> (as a percentage of total calls) from week 11 2015 - week 11 2019 (as of 20/03/2019).**



<sup>1</sup> Data supplied by Health Statistics and Analysis Unit, Welsh Government. Flu related calls are the sum of calls recorded as 'cold/flu', 'cough', 'headache', 'fever' and 'sore throat'. Following changes to the NHS Direct calls system, including the start of the 111 pilot, there has been a change in the way in which denominator data are calculated for this chart, NHS Direct Wales now count the total number of nurse triaged calls (ie calls which could have symptom data recorded against them), note that 111 includes out-of-hours calls.

## Influenza Vaccine Uptake in Wales

**Table 2. Uptake of influenza immunisations in GP Practice patients, school children and NHS staff in Wales 2018/19 (as of 20/03/2019).**

Influenza immunisation uptake in the 2018/19 season	
People aged 65y and older	68.2%
People younger than 65y in a clinical risk group	43.8%
Children aged two & three years	49.3%
Children aged four to ten years*	69.8%
NHS staff	52.3%
NHS staff who have direct patient contact	54.0%

\* In school sessions carried out so far.

The end of season report Influenza in Wales 2017/18 is available to download and contains a full breakdown of vaccination uptake amongst eligible groups.

Link to report: <http://www.wales.nhs.uk/sites3/page.cfm?orgid=457&pid=55714>

## Key points – Influenza activity in the UK and Europe

- As of week 10, influenza continued to circulate in the community with activity indicators decreasing and below baseline in the UK. Influenza GP consultations increased slightly in Scotland to 10.5 per 100,000 and decreased in Northern Ireland to 5.9 per 100,000, and remain below baseline activity in both countries. The weekly ILI GP consultation rate in England reported through the RCGP system decreased to 9.1 per 100,000 and remains below the MEM threshold for baseline activity (13.1 per 100,000). The weekly ILI consultation rate through the GP In Hours Syndromic Surveillance system was at 6.9 per 100,000 during week 10.
- During week 10, 14 samples tested positive for influenza (seven influenza A(H1N1)pdm09) and seven influenza A(H3)) through UK GP sentinel swabbing schemes. Of the 2,314 respiratory test results reported through Public Health England's DataMart scheme, there were 265 (11.5%) positive for influenza (36 influenza A(H1N1)pdm09, 111 influenza A(H3), 115 influenza A(unknown subtype) and three influenza B). UK summary data are available from the [Public Health England National Influenza Report](#).
- The WHO and the European Centre for Disease Prevention and Control (ECDC) reported that as of week 10, influenza activity is widespread throughout the WHO European Region. During week 10, a total of 1,836 sentinel specimens were tested for influenza, 786 (42.8%) of which were positive (189 influenza A(H1N1)pdm09, 236 influenza A(H3N2), 349 influenza A not subtyped and 12 influenza B). For more information on European level influenza surveillance see Flu News Europe: <http://www.flunewseurope.org/>

## World update

- The WHO reported on 18/03/19 that in the temperate zones of the northern hemisphere, influenza activity continued to be reported. In the temperate zones of the southern hemisphere, influenza activity remained at inter-seasonal levels. Worldwide, seasonal influenza subtype A accounted for the majority of influenza detections.
- Based on FluNet reporting (as of 15/03/2019), during the time period from 18/02/19 – 03/03/19, National Influenza Centres and other national influenza laboratories from 114 countries, areas or territories reported influenza surveillance data. The WHO Global Influenza Surveillance and Response System laboratories tested more than 205,150 specimens during that time period, 59,350 were positive for influenza viruses, of which 57,635 were typed as influenza A (14,751 influenza A(H1N1)pdm09, 10,037 influenza A(H3N2) and 32,847 influenza A(not subtyped)) and 1,715 influenza B (of the characterised influenza B viruses 147 belonged to the B-Yamagata lineage and 625 to the B-Victoria lineage).

**Source:** WHO influenza update:

[http://www.who.int/influenza/surveillance\\_monitoring/updates/en/](http://www.who.int/influenza/surveillance_monitoring/updates/en/)

## Update on influenza activity in North America

- The USA Centers for Disease Control and Prevention (CDC) report that during week 10 (ending 09/03/19) influenza activity decreased slightly, but remains elevated in the United States. Nationally, 124,283 (15.4%) out of 809,094 specimens have tested positive for influenza since week 40, of these positives 119,654 (96.3%) were influenza A and 4,629 (3.7%) were influenza B. Further characterisation has been carried out on 54,387 specimens by public health laboratories, and 27,552 tested positive for influenza, 26,818 (97.3%) were influenza A (17,169 influenza A(H1N1)pdm09 (67.2%), 8,382 influenza (H3N2) (32.8%), and subtyping was not performed on 1,267 specimens) and 734 influenza B (2.7%).
- The Public Health Agency of Canada reported that during week 10, influenza activity remained similar, or increased slightly, compared to the previous week. During week 10 the percentage of visits to healthcare professionals that were due to ILI was 1.3%, this is low compared to previous seasons. The percentage of tests positive for influenza increased slightly to 21.3% in week 10.

**Source:** Public Health Agency of Canada

<https://www.canada.ca/en/public-health/services/diseases/flu-influenza/influenza-surveillance/weekly-influenza-reports.html>

## **Middle East respiratory syndrome coronavirus (MERS-CoV) – latest update from WHO and ECDC**

- On 04/03/19 WHO reported eight additional cases of Middle East Respiratory Syndrome coronavirus (MERS-CoV) in Oman. Globally, 2,357 laboratory confirmed cases of human infection with MERS-CoV, including 820 associated deaths, have officially been reported to WHO since September 2012. Source: WHO Global Alert and Response website: <http://www.who.int/csr/don/archive/year/2019/en/>
- The majority of the MERS cases continue to be reported from the Middle East, and specifically from Saudi Arabia. Rapid risk assessments of the situation from ECDC, which contain epidemiological updates and advice for travellers and healthcare workers, are available from: <https://ecdc.europa.eu/en/middle-east-respiratory-syndrome-coronavirus>
- Further updates and advice for healthcare workers and travellers are available from WHO: <http://www.who.int/emergencies/mers-cov/en/> and from NaTHNaC: <https://travelhealthpro.org.uk/news/237/mers-cov-update-travelhealthpro-country-pages>

## **Human infection with avian influenza A(H7N9), China – latest update from WHO**

- The latest WHO Influenza at Human-Animal Interface summary (22/01/2019 to 12/02/2019) reports that no new cases of avian influenza A(H7N9) were reported and the risk assessment has not changed. Since February 2013, a total of 1,567 laboratory-confirmed cases of human infection with avian influenza A(H7N9), including at least 615 deaths, have been reported: [http://www.who.int/influenza/human\\_animal\\_interface/HAI\\_Risk\\_Assessment/en/](http://www.who.int/influenza/human_animal_interface/HAI_Risk_Assessment/en/)  
[http://www.fao.org/ag/againfo/programmes/en/empres/H7N9/Situation\\_update.html](http://www.fao.org/ag/againfo/programmes/en/empres/H7N9/Situation_update.html)
- The risk of international spread of avian influenza A(H7N9) is considered to be low at present. However, it is important that clinicians are aware of the possibility of human infection with animal influenza, in persons presenting with severe acute respiratory disease, while travelling or soon after returning from an area where avian influenza is a concern. Updates are available from the WHO Global Alert and Response website: <http://www.who.int/csr/don/en/>

### **Links:**

Public Health Wales influenza surveillance webpage:

<http://www.publichealthwales.org/flu-activity>

GP Sentinel Surveillance of Infections Scheme:

<http://www.wales.nhs.uk/sites3/page.cfm?orgid=457&pid=27918>

NICE influenza antiviral usage guidance:

<http://www.nice.org.uk/Guidance/TA158>

Wales influenza information:

<http://www.wales.nhs.uk/sitesplus/888/page/43745>

England influenza surveillance:

<https://www.gov.uk/government/collections/seasonal-influenza-guidance-data-and-analysis>

Scotland influenza surveillance:

<http://www.hps.scot.nhs.uk/resp/seasonalinfluenza.aspx>

Northern Ireland influenza surveillance:

<http://www.publichealth.hscni.net/directorate-public-health/health-protection/influenza>

European Centre for Communicable Disease:

<http://ecdc.europa.eu/>

European influenza information:

<http://flunewseurope.org/>

Advice on influenza immunisation (for NHS Wales users)

<http://nww.immunisation.wales.nhs.uk/home>

For further information on this report, please email Public Health Wales using:

[surveillance.requests@wales.nhs.uk](mailto:surveillance.requests@wales.nhs.uk)