

Current level of influenza activity: *Baseline activity*

Influenza activity trend: *Stable*

Confirmed influenza cases since 2020 week 40: 9 (four influenza A(H3N2), one influenza A(not subtyped) and four influenza B.)

## Key points – Wales


### Surveillance indicators suggest that influenza is not currently circulating in Wales.

The sentinel GP consultation rate for influenza-like illness (ILI) remained stable during week 46 (ending 15/11/2020). During week 46, two cases of influenza were confirmed. COVID-19 cases continue to be detected in symptomatic patients in hospital and in the community. Rhinovirus was the most commonly detected non-COVID-19 cause of Acute Respiratory Infection (ARI) but other causes of ARI continue to be detected. Surveillance data suggests the Respiratory Syncytial Virus (RSV) season is not yet underway.

- The **Sentinel GP consultation rate for influenza-like illness (ILI)** in Wales during week 46 was 0.5 consultations per 100,000 practice population (Table 1). This has remained stable compared to the previous week (0.5 consultations per 100,000) and remains below baseline threshold for seasonal influenza activity (11.0 per 100,000 practice population) (Figure 1). The sentinel GP ILI consultation rate was highest in patients aged 45-64 years (1.9 per 100,000 practice population) (Table 1). Caution should be used when comparing consultation rates from March 2020 onwards to previous periods due to the changes in health-seeking behaviours brought about by the COVID-19 pandemic.
- The **Sentinel GP consultation rate for Acute Respiratory Infections (ARI)** was 61.7 per 100,000 practice population during week 46, this is a decrease compared to the previous week (68.3 per 100,000) (Table 2). Weekly consultations for Upper Respiratory Tract Infections and Lower Respiratory Tract Infections decreased compared to the previous week. The age-group specific consultation rate for ARI during week 46 was highest in under one year olds (328.7 per 100,000 practice population).
- The percentage of calls to **NHS Direct Wales** which were 'influenza-related' (cold/flu, cough, fever, headache and sore throat) during week 46 decreased to 13.4% (Figure 9).
- The total number of respiratory-related consultations with **Out of Hours (OOH)** doctors in Wales reported to Public Health Wales during week 46 was 842, this represents 7.7% of all 10,948 reported consultations with OOH doctors, a decrease in the number and the proportion reported last week (Figure 8).
- During week 46, 256 specimens received respiratory panel testing from hospital and non-sentinel GP patients with ARI. These results do not include samples tested solely for SARS-CoV2. There was one influenza A(H3N2), one influenza B, 46 rhinovirus, 11 adenovirus and four enterovirus detected in week 46 (Figure 4). Forty-two respiratory specimens were tested from patients in intensive care units (ICU), none were positive for influenza (Figure 5). For the latest COVID-19/ SARS-CoV2 surveillance data please see the [PHW daily dashboard](#)
- Five surveillance samples from patients with ILI, collected by **sentinel GPs** during week 46 had been received by Public Health Wales Microbiology as at 18/11/2020. All samples were negative for all routinely tested respiratory pathogens.
- Surveillance data suggest that the RSV season is not yet underway with no positive samples in under 5 year olds in recent weeks.
- During week 46, 127 **ARI outbreaks** were reported to the Public Health Wales Health Protection team, 121 were reported as COVID-19 outbreaks. Sixty-five were in residential homes, 18 were in a school/nursery setting, 11 were in hospitals and 33 were in a community, mixed or other setting.
- According to **EuroMoMo** analysis, all-cause deaths are at seasonally expected levels.


## Respiratory infection activity in Wales

**Figure 1. Clinical consultation rate for ILI per 100,000 practice population in Welsh sentinel practices (as of 15/11/2020).**



\* The Moving Epidemic Method (MEM) threshold calculated for Wales ILI consultation rates is 11.1 per 100,000. MEM thresholds used in this chart are based on influenza from 2010-11 to 2018-19 seasons. Caution should be used when comparing consultation rates from March 2020 onwards to previous periods due to the changes in health-seeking behaviours brought about by the COVID-19 pandemic.

**Figure 2. Clinical consultation rate for ILI per 100,000 practice population in Welsh sentinel practices (week 48 1996 – week 46 2020).**



\* Reporting changed to Audit+ surveillance system


**Table 1. Age-specific consultations (per 100,000) for ILI in Welsh sentinel practices, week 41 – week 46 2020 (as of 15/11/2020).**

Age group	41	42	43	44	45	46
< 1	0.0	0.0	0.0	0.0	0.0	0.0
1 - 4	0.0	0.0	0.0	0.0	0.0	0.0
5 - 14	0.0	0.0	0.0	0.0	0.0	0.0
15 - 24	2.2	0.0	0.0	0.0	0.0	0.0
25 - 34	0.0	0.0	4.1	2.0	0.0	0.0
35 - 44	2.1	0.0	0.0	0.0	0.0	0.0
45 - 64	0.9	1.9	1.0	0.9	1.9	1.9
65 - 74	0.0	4.3	0.0	0.0	0.0	0.0
75+	2.4	0.0	0.0	0.0	0.0	0.0
<b>Total</b>	<b>1.0</b>	<b>1.0</b>	<b>0.8</b>	<b>0.5</b>	<b>0.5</b>	<b>0.5</b>

**Table 2. Age-specific consultations (per 100,000) for ARI in Welsh sentinel practices, week 41 – week 46 2020 (as of 15/11/2020).**


Age group	41	42	43	44	45	46
< 1	254.2	476.6	488.0	509.7	382.3	328.7
1 - 4	181.9	168.9	227.2	175.8	130.2	141.2
5 - 14	54.8	61.7	47.0	29.7	25.1	40.2
15 - 24	67.1	88.8	50.8	54.2	56.4	44.6
25 - 34	47.7	57.6	81.5	59.6	55.6	57.1
35 - 44	66.0	72.2	61.3	43.3	76.3	32.0
45 - 64	68.7	57.6	65.6	59.5	73.4	66.2
65 - 74	60.4	51.8	70.5	28.0	62.5	64.3
75+	82.1	43.4	71.7	96.4	79.5	79.3
<b>Total</b>	<b>70.3</b>	<b>68.8</b>	<b>74.0</b>	<b>61.8</b>	<b>68.3</b>	<b>61.7</b>

**Figure 3. Specimens submitted for virological testing by sentinel GPs as of 15/11/2020, by week of sample collection, week 46 2019 - week 46 2020.**




\* Tested negative for influenza, adenovirus, rhinovirus, RSV, parainfluenza, mycoplasma, human metapneumovirus, enterovirus, bocavirus and coronaviruses.

**Figure 4. Specimens submitted for virological testing for hospital patients and non-sentinel GPs as of 15/11/2020 by week of sample collection, week 46 2019 to week 46 2020.**




This chart summarises respiratory panel test data and does not include data for patients tested SOLEY for SARS-CoV2. Combined data for tests carried out in Public Health Wales Microbiology: Cardiff laboratory, provided by Public Health Wales Microbiology Cardiff Specialist Virology Centre. This chart summarises individual test results, patients who are positive for multiple infections within a given week will appear multiple times.

**Figure 5. Specimens submitted for virological testing for ICU patients, by week of sample collection, week 46 2019 to week 46 2020.**



This chart summarises respiratory panel test data and does not include data for patients tested SOLEY for SARS-CoV2.


**Figure 6. RSV incidence rate per 100,000 population aged under five years, week 30 2017 to week 46 2020.**




No samples have tested positive for RSV in under 5 year olds since week 30 2020.

### Out of Hours consultations and calls to NHS Direct Wales

**Figure 7. Weekly total consultations to Out of Hours services in Wales and numbers of respiratory-related diagnoses (as of 15/11/2020).**



**Figure 8. Influenza related calls to NHS Direct Wales<sup>1</sup> (as a percentage of total calls) from week 46 2016 - week 46 2020 (as of 15/11/2020).**



<sup>1</sup> Data supplied by Health Statistics and Analysis Unit, Welsh Government.

Flu related calls are the sum of calls recorded as 'cold/flu', 'cough', 'headache', 'fever' and 'sore throat'. Following changes to the NHS Direct calls system, including the start of the 111 pilot, there has been a change in the way in which denominator data are calculated for this chart, NHS Direct Wales now count the total number of nurse triaged calls (ie calls which could have symptom data recorded against them), note that 111 includes out-of-hours calls.

## Influenza Vaccine Uptake in Wales

**Table 3. Uptake of influenza immunisations in GP Practice patients, school children and NHS staff in Wales 2020/21 (as of 10/11/2020).**

Influenza immunisation uptake in the 2020/21 season	
People aged 65y and older	66.9%
People younger than 65y in a clinical risk group	34.2%
Children aged two & three years	42.1%
Children aged four to ten years*	72.5%
NHS staff	-
NHS staff who have direct patient contact	-

\* In school sessions carried out so far.

The end of season report Influenza in Wales 2018/19 is available to download and contains a full breakdown of vaccination uptake amongst eligible groups.

Link to report: <http://www.wales.nhs.uk/sites3/page.cfm?orgid=457&pid=55714>

NHS staff influenza vaccination uptake data for the 2020-21 season will be reported later in November.

## Influenza activity – UK and international summary

- As of week 45, the majority of community and syndromic indicators for influenza decreased or remained stable in the UK. GP ILI consultations increased in Scotland to 0.8 per 100,000, and in Northern Ireland to 1.8 per 100,000, but remains well below the baseline intensity thresholds for both countries. The weekly ILI GP consultation rate in England reported through the RCGP system increased to 1.5 per 100,000, well below the MEM threshold for baseline activity (12.2 per 100,000).
- During week 45, none of the 939 respiratory test results reported through Public Health England's DataMart scheme tested positive for influenza. UK summary data are available from the [Public Health England National Influenza and COVID-19 Surveillance Report](#).
- The WHO and the European Centre for Disease Prevention and Control (ECDC) reported that as of week 45, influenza activity remained at inter-seasonal levels across the WHO European Region. During week 45, a total of 615 sentinel specimens were tested for influenza, none of which were positive.

**Source:** Flu News Europe: <http://www.flunewseurope.org/>

- The WHO reported on 09/11/2020 that globally, influenza activity remained at lower levels than expected for this time of year. In the temperate zone of the northern hemisphere, influenza activity remained below inter-seasonal levels, though sporadic influenza detections were reported in some countries. In the temperate zones of the southern hemisphere, no influenza detections were reported across countries. In the Caribbean and Central American countries, sporadic influenza detections were reported. In tropical South America, there were no influenza detections across reporting countries. In tropical Africa, influenza activity was reported in Cote d'Ivoire, Niger and Kenya. In Southern Asia, influenza activity was reported in Bangladesh and India in recent weeks. In South East Asia, influenza detections continued to be reported in Cambodia and Lao. Worldwide, of the very few detections reported, seasonal influenza A(H3N2) viruses accounted for the majority of detections.
- Based on FluNet reporting (as of 06/11/2020), during the time period from 12/10/2020 – 25/10/2020, National Influenza Centres and other national influenza laboratories from 95 countries, areas or territories reported influenza surveillance data. The WHO Global Influenza Surveillance and Response System laboratories tested more than 94,241 specimens during that time period, 140 were positive for influenza viruses, of which 80 were typed as influenza A (two influenza A(H1N1)pdm09, 35 influenza A(H3N2) and 43 influenza A(not subtyped)) and 60 influenza B (of the characterised influenza B viruses one belonged to the B-Yamagata lineage and 14 to the B-Victoria lineage).

**Source:** WHO influenza update:

[http://www.who.int/influenza/surveillance\\_monitoring/updates/en/](http://www.who.int/influenza/surveillance_monitoring/updates/en/)

## Update on influenza activity in North America

- The USA Centers for Disease Control and Prevention (CDC) report that during week 45 (ending 07/11/2020) influenza activity remains low in the United States. Nationally, 30 (0.2%) out of 15,656 specimens have tested positive for influenza in week 45, of these positives 19 (63.3%) were influenza A and 11 (36.7%) were influenza B. Further characterisation has been carried out on 19,439 specimens by public health laboratories, and six samples tested positive for influenza, four influenza A(subtyping not performed) and two influenza B.

**Source:** CDC Weekly US Influenza Surveillance Report

<http://www.cdc.gov/flu/weekly/>

- The Public Health Agency of Canada reported that during week 45, influenza activity remains below average for this time of year. The percentage of visits to healthcare professionals that were due to ILI was 0.6% in week 45, similar to previous weeks. The percentage of tests positive for influenza was 0.04% during week 45.

**Source:** Public Health Agency of Canada

<https://www.canada.ca/en/public-health/services/diseases/flu-influenza/influenza-surveillance/weekly-influenza-reports.html>

### **Coronavirus disease 2019 (COVID-19) – UK and international summary**

- The number of confirmed cases in Wales reported as at 18/11/2020 is 68,449, with 640 newly reported in the previous 24 hours. The cumulative number of suspected COVID-19 deaths in confirmed cases in hospitals and care homes reported to Public Health Wales is 2,284, with 41 new deaths reported in the previous 24 hours. The cumulative number of registered deaths in Welsh residents where COVID-19 was mentioned in the death certificate as at 2020 week 44 was 2,879. Latest COVID-19 data from Public Health Wales is available from: <https://phw.nhs.wales/topics/latest-information-on-novel-coronavirus-covid-19/> Public Health Wales produce a daily statement on COVID-19, available from: <https://covid19-phwstatement.nhs.wales/>
- As at 18/11/2020, there are 1,430,341 reported confirmed cases of COVID-19 in the UK, of which 19,609 were newly reported in the previous 24 hours. The total deaths within 28 days of a positive test was 53,274, with 529 reported in the previous 24 hours. Latest UK data is available from: [https://coronavirus.data.gov.uk/?\\_ga=2.47134183.1732144231.1599825067-744978499.1577716555](https://coronavirus.data.gov.uk/?_ga=2.47134183.1732144231.1599825067-744978499.1577716555)
- As at 18/11/2020, WHO reported 55,064,128 confirmed COVID-19 cases globally, with 291,514 reported in the previous 24 hours. There have been 1,328,015 deaths, of which 3,757 were reported in the previous 24 hours. Daily WHO situation updates are available from: <https://covid19.who.int/>

### **Middle East respiratory syndrome coronavirus (MERS-CoV) – latest update from WHO and ECDC**

- On 02/07/2020 WHO reported nine additional cases of Middle East Respiratory Syndrome coronavirus (MERS-CoV), including five associated deaths. Globally, 2,562 laboratory confirmed cases of human infection with MERS-CoV, including 881 associated deaths, have officially been reported to WHO since 2012.  
Source: WHO Global Alert and Response website: <http://www.who.int/csr/don/archive/year/2020/en/>
- The majority of the MERS cases continue to be reported from the Middle East, and specifically from Saudi Arabia. Rapid risk assessments of the situation from ECDC, which contain epidemiological updates and advice for travellers and healthcare workers, are available from: <https://ecdc.europa.eu/en/middle-east-respiratory-syndrome-coronavirus>
- Further updates and advice for healthcare workers and travellers are available from WHO: <http://www.who.int/emergencies/mers-cov/en/> and from NaTHNaC: <https://travelhealthpro.org.uk/news/237/mers-cov-update-travelhealthpro-country-pages>

### **Human infection with avian influenza A(H7N9), China – latest update from WHO**

- The latest WHO Influenza at Human-Animal Interface summary (11/07/2020 to 23/10/2020) reports that no new cases of avian influenza A(H7N9) were reported. Since February 2013, a total of 1,568 laboratory-confirmed cases of human infection with avian influenza A(H7N9), including at least 616 deaths, have been reported: [http://www.who.int/influenza/human\\_animal\\_interface/HAI\\_Risk\\_Assessment/en/](http://www.who.int/influenza/human_animal_interface/HAI_Risk_Assessment/en/) [http://www.fao.org/ag/againfo/programmes/en/empres/H7N9/Situation\\_update.html](http://www.fao.org/ag/againfo/programmes/en/empres/H7N9/Situation_update.html)
- The risk of international spread of avian influenza A(H7N9) is considered to be low at present. However, it is important that clinicians are aware of the possibility of human infection with animal influenza, in persons presenting with severe acute respiratory disease, while travelling or soon after returning from an area where avian influenza is a concern. WHO Global Alert & Response updates: <http://www.who.int/csr/don/en/>



**Links:**

**Public Health Wales influenza surveillance webpage:**

<http://www.wales.nhs.uk/sites3/page.cfm?orgid=457&pid=25480>

**Public Health Wales COVID-19 data dashboard:**

<https://public.tableau.com/profile/public.health.wales.health.protection#!/vizhome/RapidCOVID-19virology-Public/Headlinesummary>

**GP Sentinel Surveillance of Infections Scheme:**

<http://www.wales.nhs.uk/sites3/page.cfm?orgid=457&pid=27918>

**NICE influenza antiviral usage guidance:**

<http://www.nice.org.uk/Guidance/TA158>

**Wales influenza information:**

<https://phw.nhs.wales/topics/flu/>

**England influenza surveillance:**

<https://www.gov.uk/government/statistics/weekly-national-flu-reports-2019-to-2020-season>

**Scotland influenza surveillance:**

<https://www.hps.scot.nhs.uk/a-to-z-of-topics/influenza/#data>

**Northern Ireland influenza surveillance:**

<https://www.publichealth.hscni.net/directorate-public-health/health-protection/seasonal-influenza>

**European Centre for Communicable Disease:**

<http://ecdc.europa.eu/>

**European influenza information:**

<http://flunewseurope.org/>

**Advice on influenza immunisation (for NHS Wales users)**

<http://www.immunisation.wales.nhs.uk/home>

For further information on this report, please email Public Health Wales using:

[surveillance.requests@wales.nhs.uk](mailto:surveillance.requests@wales.nhs.uk)