

lechyd Cyhoeddus Cymru Public Health Wales

Current level of influenza activity: *Baseline activity* Influenza activity trend: *Stable* Confirmed influenza cases since 2020 week 40: 38 (10 influenza A(H3N2), 15 influenza A(not subtyped) and 13 influenza B).

Key points – Wales


Surveillance indicators suggest that RSV is circulating in Wales and influenza is not.

During week 27 (ending 11/07/2021) there has been a continued increase in numbers of confirmed cases of Respiratory Synctial Virus (RSV) in children aged under 5 years, mainly in north Wales, and this week activity remains above the threshold that would normally indicate seasonal circulation. RSV did not circulate over the 2020-21 winter. The current increase in cases is earlier than the usual RSV season in Wales and it is unclear whether it will follow the usual epidemic pattern for RSV, there is potential for higher than usual numbers of cases this year. No influenza cases were confirmed during week 27. COVID-19 cases continue to be detected in symptomatic patients in hospital and in the community with continuing increases in recent weeks. Rhinovirus and parainfluenza are the most commonly detected cause of non-COVID-19 Acute Respiratory Infection (ARI), confirmed cases of parainfluenza (type 3) have increased significantly recently, particularly in children.

- The Sentinel GP consultation rate for influenza-like illness (ILI) in Wales during week 27 was 0.3 consultations per 100,000 practice population (Table 1). This remains stable compared to the previous week (0.3 consultations per 100,000) and remains well below baseline threshold for seasonal influenza activity (11.0 per 100,000 practice population) (Figure 1). Caution should be used when comparing consultation rates from March 2020 onwards to previous periods due to the changes in health-seeking behaviours brought about by the COVID-19 pandemic.The Sentinel GP consultation rate for Acute Respitatory Infections (ARI) was 50.8 per 100,000 practice population during week 27, this is a decrease compared to the previous week (57.0 per 100,000) (Table 2). Weekly consultations for Upper Respiratory Tract Infections and Lower Respiratory Tract Infections decreased compared to the previous week. The age-group specific consultation rate for ARI during week 27 was highest in under one year olds (684.4 per 100,000 practice population). In recent weeks ARI consultations have increased in children aged under 5 years.
- The percentage of calls to **NHS Direct Wales** which were 'influenza-related' (cold/flu, cough, fever, headache and sore throat) during week 26 decreased to 19.3% (Figure 8).
- During week 27, 1,354 specimens received respiratory panel testing, mainly from patients attending hospitals with ARI symptoms. These results do not include samples tested solely for SARS-CoV2. There were 19 RSV, 208 parainfluenza, 176 rhinoviruses, 35 adenoviruses, 23 enteroviruses, three seasonal coronaviruses, two human metapneumoviruses and one bocavirus, detected in week 27 (Figure 4). Additionally, 4,119 samples from patients were tested for influenza, RSV and SARS-CoV2 only, many of these tests may be associated with screening activities rather than diagnostic testing for patients presenting with ARI symptoms. Of these 4,119 samples, none were positive for influenza, nine were positive for RSV and 117 were positive for SARS-CoV2 (Figure 5). Thirty-eight respiratory specimens were tested from patients in intensive care units (ICU) and none were positive for influenza (Figure 6). For the latest COVID-19/ SARS-CoV2 surveillance data please see the PHW daily dashboard
- No surveillance samples from patients with ILI, collected by **sentinel GPs** during week 27, had been received by Public Health Wales Microbiology as at 14/07/2021.
- Confirmed RSV case numbers in children aged under 5 continue to increase. Activity has exceeded the threshold that would usually indicate onset of circulation. In week 27 there were 12.9 confirmed cases per 100,000 in this age group (Figure 7). The provisional MEM threshold in Wales which predicts the start of the annual RSV season in children younger than five years is 6.3 confirmed cases per 100,000.
- During week 27, 55 **ARI outbreaks** were reported to the Public Health Wales Health Protection team, all were reported as COVID-19 outbreaks. Twenty-eight were in residential homes, eight were in a school/nursery setting, one was in a hospital and eighteen were in a community, mixed or other setting.
- According to **EuroMoMo** analysis, all-cause deaths were not significantly in excess during week 26 (latest data).

Respiratory infection activity in Wales

Figure 1. Clinical consultation rate for ILI per 100,000 practice population in Welsh sentinel practices (as of 11/07/2021).



* The Moving Epidemic Method (MEM) threshold calculated for Wales ILI consultation rates is 11.1 per 100,000. MEM thresholds used in this chart are based on influenza from 2010-11 to 2018-19 seasons. Caution should be used when comparing consultation rates from March 2020 onwards to previous periods due to the changes in health-seeking behaviours brought about by the COVID-19 pandemic.




Figure 2. Clinical consultation rate for ILI per 100,000 practice population in Welsh

Reporting changed to Audit+ surveillance system


Table 1. Age-specific consultations (per 100,000) for ILI in Welsh sentinel practices, week 21– week 27 2021 (as of 11/07/2021).

Age						
group	22	23	24	25	26	27
< 1	0.0	0.0	0.0	0.0	0.0	0.0
1 - 4	0.0	0.0	0.0	0.0	0.0	0.0
5 - 14	0.0	0.0	0.0	0.0	0.0	0.0
15 - 24	0.0	2.2	0.0	0.0	0.0	2.4
25 - 34	2.0	0.0	0.0	0.0	0.0	0.0
35 - 44	2.0	4.0	0.0	0.0	2.0	0.0
45 - 64	0.0	0.0	1.0	0.0	0.0	0.0
65 - 74	0.0	0.0	0.0	2.2	0.0	0.0
75+	0.0	0.0	0.0	0.0	0.0	0.0
Total	0.5	0.7	0.3	0.3	0.3	0.3

Table 2. Age-specific consultations (per 100,000) for ARI in Welsh sentinel practices, week 21 – week 27 2021 (as of 11/07/2021).


Age						
group	22	23	24	25	26	27
< 1	367.3	758.6	977.8	1342.3	665.8	684.4
1 - 4	455.4	614.2	768.1	942.3	500.5	430.1
5 - 14	36.7	65.8	76.4	94.8	55.1	47.1
15 - 24	84.7	70.8	56.4	79.3	38.2	41.2
25 - 34	53.7	70.7	56.2	62.0	45.6	38.2
35 - 44	36.9	64.7	43.4	61.7	30.6	30.8
45 - 64	39.6	56.5	41.0	53.3	24.4	23.5
65 - 74	30.7	45.1	55.9	57.5	30.4	19.1
75+	51.7	61.7	45.1	39.0	33.5	24.0
Total	64.1	87.3	85.5	105.0	57.0	50.8

Figure 3. Specimens submitted for virological testing by sentinel GPs as of 11/07/2021, by week of sample collection, week 28 2020 to week 27 2021.




* Tested negative for influenza, adenovirus, rhinovirus, RSV, parainfluenza, mycoplasma, human metapneumovirus, enterovirus, bocavirus and coronaviruses.

Figure 4. Specimens submitted for virological testing for hospital patients and non-sentinel GPs as of 14/07/2021 by week of sample collection, week 28 2020 to week 27 2021.




This chart summarises respiratory panel test data and does not include data for patients tested SOLEY for SARS-CoV2. Combined data for tests carried out in Public Health Wales Microbiology: Cardiff laboratory, provided by Public Health Wales Microbiology Cardiff Specialist Virology Centre. This chart summarises individual test results, patients who are positive for multiple infections within a given week will appear multiple times.











This chart summarises respiratory panel test data and does not include data for patients tested SOLEY for SARS-CoV2.

Figure 7. RSV incidence rate per 100,000 population aged under five years, week 31 2017 to week 27 2021.





¹ Data supplied by Health Statistics and Analysis Unit, Welsh Government. Flu related calls are the sum of calls recorded as 'cold/flu', 'cough', 'headache', 'fever' and 'sore throat'. Following changes to the NHS Direct calls system, including the start of the 111 pilot, there has been a change in the way in which denominator data are calculated for this chart, NHS Direct Wales now count the total number of nurse triaged calls (ie calls which could have symptom data recorded against them), note that 111 includes out-of-hours calls.

Influenza Vaccine Uptake in Wales

Table 3. Uptake of influenza immunisations in GP Practice patients, school children andNHS staff in Wales 2020/21 (as of 23/03/2021).

Influenza immunisation uptake in the 2020/21 season					
People aged 65y and older	76.5%				
People younger than 65y in a clinical risk group	51.0%				
Children aged two & three years	56.3%				
Children aged four to ten years*	72.4%				
NHS staff	63.4%				
NHS staff who have direct patient contact	65.2%				

* In school sessions carried out so far.

The end of season report Influenza in Wales 2019/20 is available to download and contains a full breakdown of vaccination uptake amongst eligible groups. Link to report: http://www.wales.nhs.uk/sites3/page.cfm?orgid=457&pid=55714

Influenza activity - UK and international summary

- As of week 26, the majority of community and syndromic influenza indicators remained low in the UK. GP ILI consultations remained stable in Scotland at 0.5 per 100,000; Northern Ireland was at 0.6 per 100,000 (as at week 25), well below the baseline intensity thresholds. The weekly ILI GP consultation rate in England reported through the RCGP system decreased slightly to 0.9 per 100,000, remaining well below the MEM threshold for baseline activity (12.2 per 100,000).
- During week 26, one of the 2,934 respiratory test results reported through Public Health England's
 DataMart scheme tested positive for influenza. UK summary data are available from the <u>Public Health</u>
 <u>England National Influenza and COVID-19 Surveillance Report</u>.
- The WHO and the European Centre for Disease Prevention and Control (ECDC) reported that during weeks 21-24, influenza activity remained at inter-seasonal levels across the WHO European Region.
 Source: Flu News Europe: <u>http://www.flunewseurope.org/</u>
- The WHO reported on 05/07/2021 that globally, despite continued or even increased testing in some countries, influenza activity remained at lower levels than expected for this time of year. In the temperate zone of the northern hemisphere, influenza activity remained at inter-seasonal levels, though detections of influenza B-Victoria lineage slightly increased, especially in China. In the temperate zones of the southern hemisphere, influenza activity remained at inter-seasonal levels. In the Caribbean and Central American countries, there were no influenza detections reported. In tropical South America, no influenza detections were reported. In tropical Africa, a few influenza detections were reported in some countries in Western and Eastern Africa. In Southern Asia, a few influenza detections were reported from India. In South East Asia, no influenza detections were reported. Worldwide, influenza B accounted for the majority of the very low numbers of detections reported.
- Based on FluNet reporting (as of 02/07/2021), during the time period from 07/06/2021 20/06/2021, National Influenza Centres and other national influenza laboratories from 75 countries, areas or territories reported influenza surveillance data. The WHO Global Influenza Surveillance and Response System laboratories tested more than 146,816 specimens during that time period, 766 were positive for influenza viruses, of which 84 were typed as influenza A (of the subtyped influenza A viruses, 48 were influenza A(H1N1)pdm09 and eight were influenza A(H3N2)) and 682 influenza B (of the characterised influenza B viruses 624 belonged to the B-Victoria lineage).

Source: WHO influenza update:

https://www.who.int/teams/global-influenza-programme/surveillance-and-monitoring/influenzaupdates/current-influenza-update

Australia and New Zealand update

In New Zealand, during the week ending 04/07/2021, the GP influenza-like illness activity (ILI) consultation rates have continued to increase. There have been no influenza viruses detected this season to date, rhinovirus has been the most commonly detected virus. RSV was the virus most commonly detected through SARI surveillance in the past week, with high numbers of hospitalisations across the country.

Source: Institute of Environmental Science & Research, New Zealand https://www.esr.cri.nz/our-services/consultancy/flu-surveillance-and-research

• In Australia, according to the latest available update (fortnight ending 04/07/2021), influenza-like illness (ILI) activity in the community remains at low levels, but has continued to gradually increase since January 2021. To date, the majority of nationally reported laboratory-confirmed influenza cases were influenza A (70.4%).

Source: Australian Influenza Surveillance Report and Activity Updates. https://www1.health.gov.au/internet/main/publishing.nsf/Content/cda-surveil-ozflu-flucurr.htm#current

Respiratory syncytial virus (RSV) in North America

• The USA CDC has reported an out of season increase in RSV activity, with an increase in sample positivity since early March 2021.

Source: CDC RSV national trends

https://www.cdc.gov/surveillance/nrevss/rsv/natl-trend.html

Coronavirus disease 2019 (COVID-19) – UK and international summary

- The number of confirmed cases in Wales reported as at 14/07/2021 is 227,563, with 1,135 newly reported in the previous 24 hours. The cumulative number of suspected COVID-19 deaths in confirmed cases in hospitals and care homes reported to Public Health Wales is 5,581, with one new death reported in the previous 24 hours. The cumulative number of registered deaths in Welsh residents where COVID-19 was mentioned in the death certificate as at 2021 week 25 was 7,893. Latest COVID-19 data from Public Health Wales is available from: https://phw.nhs.wales/topics/latest-information-on-novel-coronavirus-covid-19/ Public Health Wales produce a daily statement on COVID-19, available from: https://covid19-phwstatement.nhs.wales/
- As at 13/07/2021, there have been 5,191,459 reported confirmed cases of COVID-19 in the UK, of which 36,660 were newly reported in the previous 24 hours. The total deaths within 28 days of a positive test was 128,481, with 50 reported in the previous 24 hours. Latest UK data is available from: <u>https://coronavirus.data.gov.uk/</u>
- As at 07/07/2021, WHO have reported 187,086,096 confirmed COVID-19 cases globally, with 372,970 reported in the previous 24 hours. There have been 4,042,921 deaths, of which 7,469 were reported in the previous 24 hours. Daily WHO situation updates are available from: https://covid19.who.int/

Middle East respiratory syndrome coronavirus (MERS-CoV) – latest update from WHO and ECDC

 On 14/04/2021 WHO reported seven additional cases of Middle East Respiratory Syndrome coronavirus (MERS-CoV), including three associated deaths. Globally, 2,574 laboratory confirmed cases of human infection with MERS-CoV, including 886 associated deaths, have officially been reported to WHO since 2012.

Source: WHO Global Alert and Response website: <u>https://www.who.int/emergencies/disease-outbreak-news</u>

- The majority of the MERS cases continue to be reported from the Middle East, and specifically from Saudi Arabia. Rapid risk assessments of the situation from ECDC, which contain epidemiological updates and advice for travellers and healthcare workers, are available from: <u>https://ecdc.europa.eu/en/middle-eastrespiratory-syndrome-coronavirus</u>
- Further updates and advice for healthcare workers and travellers are available from WHO: <u>http://www.who.int/emergencies/mers-cov/en/</u> and from NaTHNaC: <u>https://travelhealthpro.org.uk/news/237/mers-cov-update-travelhealthpro-country-pages</u>

Human infection with avian influenza A(H7N9), China

• The latest WHO Influenza at Human-Animal Interface summary (16/04/2021 to 21/05/2021) reports that no new cases of avian influenza A(H7N9) were reported. Since February 2013, a total of 1,568 laboratory-confirmed cases of human infection with avian influenza A(H7N9), including at least 616 deaths, have been reported:

https://www.who.int/publications/m/item/influenza-at-the-human-animal-interface-summary-and-assessment-21may-2021

http://www.fao.org/ag/againfo/programmes/en/empres/H7N9/Situation_update.html

The risk of international spread of avian influenza A(H7N9) is considered to be low at present. However, it
is important that clinicians are aware of the possibility of human infection with animal influenza, in persons
presenting with severe acute respiratory disease, while travelling or soon after returning from an area
where avian influenza is a concern. WHO Global Alert & Response updates:
https://www.who.int/emergencies/disease-outbreak-news

Links:

Public Health Wales influenza surveillance webpage: http://www.wales.nhs.uk/sites3/page.cfm?orgid=457&pid=25480 Public Health Wales COVID-19 data dashboard: https://public.tableau.com/profile/public.health.wales.health.protection#!/vizhome/RapidCOVID-19virology-Public/Headlinesummary **GP Sentinel Surveillance of Infections Scheme:** http://www.wales.nhs.uk/sites3/page.cfm?orgid=457&pid=27918 NICE influenza antiviral usage guidance: http://www.nice.org.uk/Guidance/TA158 Wales influenza information: https://phw.nhs.wales/topics/flu/ England influenza and COVID-19 surveillance: https://www.gov.uk/government/statistics/national-flu-and-covid-19-surveillance-reports Scotland seasonal respiratory surveillance: https://beta.isdscotland.org/find-publications-and-data/population-health/covid-19/weekly-national-seasonalrespiratory-report/ Northern Ireland influenza surveillance: https://www.publichealth.hscni.net/directorate-public-health/health-protection/seasonal-influenza **European Centre for Communicable Disease:** http://ecdc.europa.eu/ European influenza information: http://flunewseurope.org/ Advice on influenza immunisation (for NHS Wales users) http://nww.immunisation.wales.nhs.uk/home

For further information on this report, please email Public Health Wales using: surveillance.requests@wales.nhs.uk