# Public Health Wales CDSC Weekly Influenza & Acute Respiratory Infection Surveillance Report



Wednesday 9th March 2022 (covering week 9 2022)

Current level of influenza activity: Baseline activity

Influenza activity trend: Stable

Confirmed influenza cases since 2021 week 40: 222 (56 influenza A(H3N2), four influenza A(H1N1)pdm09, 118

influenza A(not subtyped) and 44 influenza B).

#### **Key points - Wales**


<u>Surveillance indicators suggest that although influenza is not circulating widely, confirmed case numbers</u> have increased in recent weeks.

During Week 09 (ending 06/03/2022) there were 18 cases of influenza confirmed. Confirmed cases of Respiratory Synctial Virus (RSV) in children aged under 5 years are at baseline levels. COVID-19 cases continue to be detected in symptomatic patients in hospital and in the community. Rhinovirus and adenovirus are the most commonly detected cause of non-COVID-19 Acute Respiratory Infection (ARI), with increasing confirmed cases in recent weeks.

- The Sentinel GP consultation rate for influenza-like illness (ILI) in Wales during week 09 was 2.6 consultations per 100,000 practice population (Table 1). This increased compared to the previous week (1.2 consultations per 100,000) but remains well below baseline threshold for seasonal influenza activity (11.0 per 100,000 practice population) (Figure 1). Caution should be used when comparing consultation rates from March 2020 onwards to previous periods due to the changes in health-seeking behaviours brought about by the COVID-19 pandemic.
- The Sentinel GP consultation rate for Acute Respitatory Infections (ARI) was 129.4 per 100,00 practice
  population during Week 09, this is an increase compared to the previous week (125.5 per 100,000) (Table 2).
  Weekly consultations for Upper Respiratory Tract Infections and Lower Respiratory Tract Infections increased
  compared to the previous week.
- The percentage of calls to **NHS Direct Wales** which were 'influenza-related' (cold/flu, cough, fever, headache and sore throat) during Week 09 increased to 16.2% (Figure 8).
- During Week 09, 1,755 specimens received multiplex respiratory panel testing mainly from patients attending hospitals. These results do not include samples tested solely for SARS-CoV2. There were eight influenza (five influenza A(H3N2), two influenza A(not typed) and one influenza B), eight RSV, 175 SARS-CoV2, 211 rhinoviruses, 60 adenoviruses, 32 enteroviruses, 20 parainfluenza and six human metapneumoviruses detected in Week 09 (Figure 4). Additionally, 4,124 samples from patients were tested for influenza, RSV and SARS-CoV2 only, many of these tests may be associated with screening activities rather than diagnostic testing for patients presenting with ARI symptoms. Of these 4,124 samples, 10 were positive for influenza (all untyped influenza A), four were positive for RSV and 266 were positive for SARS-CoV2 (Figure 5). Eighty-eight respiratory specimens were tested from patients in intensive care units (ICU) and none were positive for influenza (Figure 6). For the latest COVID-19/ SARS-CoV2 surveillance data please see the PHW daily dashboard
- There were two surveillance samples from patients with ILI collected by **sentinel GPs** during Week 09 (as at 09/03/2022), one sample was positive for parainfluenza and the other sample was negative for all routinely tested respiratory pathogens.
- Confirmed RSV case incidence in children aged under 5 increased, but remains at the threshold that would usually indicate baseline levels of circulation. In week 09 there were 4.1 confirmed cases per 100,000 in this age group (Figure 7). The provisional MEM threshold in Wales which predicts the start of the annual RSV season in children younger than five years is 6.3 confirmed cases per 100,000.
- During Week 09, 36 ARI outbreaks were reported to the Public Health Wales Health Protection team, all were reported as COVID-19 outbreaks. 32 were in residential homes and four were in a community, mixed or other setting.
- According to EuroMoMo analysis, all-cause deaths in Wales were not significantly in excess during week 08.


# Respiratory infection activity in Wales

Figure 1. Clinical consultation rate for ILI per 100,000 practice population in Welsh sentinel practices (as of 06/03/2022).



<sup>\*</sup> The Moving Epidemic Method (MEM) threshold calculated for Wales ILI consultation rates is 11.1 per 100,000. MEM thresholds used in this chart are based on influenza from 2010-11 to 2018-19 seasons. Caution should be used when comparing consultation rates from March 2020 onwards to previous periods due to the changes in health-seeking behaviours brought about by the COVID-19 pandemic.

Figure 2. Clinical consultation rate for ILI per 100,000 practice population in Welsh sentinel practices (week 48 1996 – week 09 2022).



<sup>\*</sup> Reporting changed to Audit+ surveillance system


Table 1. Age-specific consultations (per 100,000) for ILI in Welsh sentinel practices, week 04 – week 09 2022 (as of 06/03/2022).

Age						
group	4	5	6	7	8	9
< 1	0.0	0.0	0.0	0.0	0.0	0.0
1 - 4	6.7	0.0	0.0	0.0	0.0	0.0
5 - 14	0.0	0.0	0.0	0.0	0.0	0.0
15 - 24	0.0	2.2	2.2	0.0	0.0	2.6
25 - 34	0.0	2.0	2.0	0.0	5.8	0.0
35 - 44	0.0	2.0	4.0	2.0	0.0	4.7
45 - 64	1.9	0.9	2.8	1.9	0.9	4.4
65 - 74	0.0	2.2	0.0	0.0	0.0	5.2
75+	0.0	4.7	0.0	0.0	2.3	0.0
Total	0.7	1.7	1.8	0.8	1.2	2.6

Table 2. Age-specific consultations (per 100,000) for ARI in Welsh sentinel practices, week 04 – week 09 2022 (as of 06/03/2022).


Age						
group	4	5	6	7	8	9
< 1	501.6	663.7	768.3	486.4	486.4	607.2
1 - 4	477.0	564.2	610.6	485.1	499.0	427.3
5 - 14	149.4	123.7	126.2	142.4	123.7	118.9
15 - 24	103.7	134.0	142.9	113.0	117.5	131.7
25 - 34	126.5	135.7	129.9	117.9	123.9	126.5
35 - 44	123.1	100.4	120.6	102.0	91.8	122.8
45 - 64	77.6	71.2	100.4	112.6	105.0	108.7
65 - 74	71.5	72.4	80.7	93.2	91.0	113.6
75+	119.7	93.1	102.5	87.6	111.3	80.5
Total	121.8	120.8	136.6	126.8	125.5	129.4

Figure 3. Specimens submitted for virological testing by sentinel GPs as of 06/03/2022, by week of sample collection, week 09 2021 to week 09 2022.



<sup>\*</sup> Tested negative for influenza, adenovirus, rhinovirus, RSV, parainfluenza, mycoplasma, human metapneumovirus, enterovirus, bocavirus and coronaviruses. Samples which test positive for more than on pathogen will appear more than once in the chart.

Figure 4. Specimens submitted for virological testing for hospital patients and non-sentinel GPs as of 06/03/2022 by week of sample collection, week 09 2021 to week 09 2022.



This chart summarises respiratory panel test data and does not include data for patients tested SOLEY for SARS-CoV2. Combined data for tests carried out in Public Health Wales Microbiology: Cardiff laboratory, provided by Public Health Wales Microbiology Cardiff Specialist Virology Centre. This chart summarises individual test results, patients who are positive for multiple infections within a given week will appear multiple times. Samples which test positive for more than on pathogen will appear more than once in the chart.

Figure 5. Specimens from hospital patients submitted for RSV, Influenza and SARS-CoV2 testing only, as of 06/03/2022 by week of sample collection, week 09 2021 to week 09 2022.






Figure 6. Specimens submitted for virological testing for ICU patients, by week of sample collection, week 09 2021 to Week 09 2022.




This chart summarises respiratory panel test data and does not include data for patients tested SOLEY for SARS-CoV2. Samples which test positive for more than on pathogen will appear more than once in the chart.

Figure 7. RSV incidence rate per 100,000 population aged under five years, week 30 2017 to Week 09 2022.



## Calls to NHS Direct Wales

Figure 8. Influenza related calls to NHS Direct Wales<sup>1</sup> (as a percentage of total calls) from week 10 2018 - Week 09 2022 (as of 06/03/2022).



<sup>&</sup>lt;sup>1</sup> Data supplied by Health Statistics and Analysis Unit, Welsh Government.

Flu related calls are the sum of calls recorded as 'cold/flu', 'cough', 'headache', 'fever' and 'sore throat'. Following changes to the NHS Direct calls system, including the start of the 111 pilot, there has been a change in the way in which denominator data are calculated for this chart, NHS Direct Wales now count the total number of nurse triaged calls (ie calls which could have symptom data recorded against them), note that 111 includes out-of-hours calls.

# Influenza Vaccine Uptake in Wales

Table 3. Uptake of influenza immunisations in GP Practice patients, school children and NHS staff in Wales 2021/22 (as of 03/03/2022).

Influenza immunisation uptake in the 2021/22 season				
People aged 65y and older	77.8%			
People younger than 65y in a clinical risk group	47.9%			
Children aged two & three years	47.5%			
Children aged four to ten years*	68.2%			
Children aged 11 to 15 years*	60.3%			
NHS staff	54.8%			
NHS staff who have direct patient contact	56.0%			

<sup>\*</sup> In school sessions carried out so far.

The end of season report Influenza in Wales 2019/20 is available to download and contains a full breakdown of vaccination uptake amongst eligible groups.

Link to report: http://www.wales.nhs.uk/sites3/page.cfm?orgid=457&pid=55714

# Influenza activity – UK and international summary

- As of week 08, community and syndromic influenza indicators remain very low in the UK. GP ILI consultations increased in Scotland at 0.8 per 100,000 and decreased in Northern Ireland to 0.9 per 100,000 remaining well below the baseline intensity thresholds. The weekly ILI GP consultation rate in England reported through the RCGP system decreased to 1.0 per 100,000, well below the MEM threshold for baseline activity (12.2 per 100,000). During week 08, 61 of the 5,706 samples tested positive for influenza (including eight influenza A(H1N1)pdm09, 18 influenza A(H3N2), 34 influenza A(not subtyped) and one influenza B). UK summary data are available from the UKHSA Influenza and COVID-19 Surveillance Report.
- The WHO and the European Centre for Disease Prevention and Control (ECDC) reported that during week 08, influenza activity continues to be reported throughout the WHO European Region. During week 08, a total of 948 sentinel specimens were tested for influenza, 92 of which were positive, 91 influenza A (74 influenza A(H3), three influenza A(H1)pdm09 and 14 influenza A(not subtyped)) and one influenza B.
   Source: Flu News Europe: http://www.flunewseurope.org/
- The WHO reported on 21/02/2022 that globally, influenza activity remained low and decreased. In the temperate zones of the northern hemisphere, influenza activity decreased with detections of mainly influenza A(H3N2) and influenza B/Victoria lineage reported. In the temperate zones of the southern hemisphere, influenza activity remained low overall. In North America, influenza detections decreased. In Europe, influenza activity appeared to decrease. In East Asia, influenza activity decreased in China and remained low in the rest of the subregion. In Northern Africa, influenza activity increased. In Western Asia, influenza activity continued to decrease. In the Caribbean and Central American countries, some influenza activity was reported with influenza A(H3N2) predominating. In tropical South America, some influenza activity was reported with influenza A(H3N2) predominating. In tropical Africa, influenza activity was reported mainly from Eastern and Middle Africa. In Southern Asia, influenza virus detections of predominantly influenza A(H3N2) decreased. In South-East Asia, mainly influenza A(H3N2) detections were reported as well as some influenza B.
- Based on FluNet reporting (as of 04/02/2022), during the time period from 24/01/2022 06/02/2022, National Influenza Centres and other national influenza laboratories from 113 countries, areas or territories reported influenza surveillance data. The WHO Global Influenza Surveillance and Response System laboratories tested more than 490,516 specimens during that time period, 12,368 were positive for influenza viruses, of which 8,423 were typed as influenza A (of the subtyped influenza A viruses, 170 were influenza A(H1N1)pdm09 and 2,483 were influenza A(H3N2)) and 3,945 influenza B (of the characterised influenza B viruses four belonged to B-yamagata lineage and 3,713 belonged to the B-Victoria lineage).

**Source:** WHO influenza update: <a href="https://www.who.int/teams/global-influenza-programme/surveillance-and-monitoring/influenza-updates/current-influenza-update">https://www.who.int/teams/global-influenza-programme/surveillance-and-monitoring/influenza-updates/current-influenza-update</a>

#### Update on influenza activity in North America

- The USA Centers for Disease Control and Prevention (CDC) report that during week 08 (ending 26/02/2022) sporadic influenza activity continues across the United States. Nationally, 1,462 (4.1%) out of 35,541 specimens have tested positive for influenza in week 08, of these positives 1,445 (98.8%) were influenza A and 17 (1.2%) were influenza B. Further characterisation has been carried out on 17,331 specimens by public health laboratories, and 293 samples tested positive for influenza, 106 influenza A(H3N2), 185 influenza A(not subtyped) and two influenza B(lineage not performed).
  - Source: CDC Weekly US Influenza Surveillance Report: http://www.cdc.gov/flu/weekly/
- The Public Health Agency of Canada reported that during week 08, influenza activity has decreased in recent weeks and remains low for this time of year. The percentage of visits to healthcare professionals that were due to ILI was 0.6% in week 08. The percentage of tests positive for influenza was 0.1% during week 08.
   Source: Public Health Agency of Canada: <a href="https://www.canada.ca/en/public-health/services/diseases/flu-influenza/influenza-surveillance/weekly-influenza-reports.html">https://www.canada.ca/en/public-health/services/diseases/flu-influenza-surveillance/weekly-influenza-reports.html</a>

## Respiratory syncytial virus (RSV) in North America

• The USA CDC has reported an out of season increase in RSV activity, with an increase in sample positivity since early March 2021, but has seen a downward trend in recent weeks.

Source: CDC RSV national trends: https://www.cdc.gov/surveillance/nrevss/rsv/natl-trend.html

# **COVID-19 – UK and international summary**

- The number of confirmed cases in Wales reported as at 08/03/2022 is 821,797 with 2,628 newly reported in the previous 24 hours. The cumulative number of suspected COVID-19 deaths in confirmed cases in hospitals and care homes reported to Public Health Wales is 7,033 with 16 new deaths reported in the previous 24 hours. The cumulative number of registered deaths in Welsh residents where COVID-19 was mentioned in the death certificate as at 2022 week 07 was 9,687. Latest COVID-19 data from Public Health Wales is available from: <a href="https://public.tableau.com/app/profile/public.health.wales.health.protection/viz/RapidCOVID-19virology-Public/Headlinesummary">https://public.tableau.com/app/profile/public.health.wales.health.protection/viz/RapidCOVID-19virology-Public/Headlinesummary</a>
- As at 08/03/2022, there have been 19,307,014 reported confirmed cases of COVID-19 in the UK, of which 322,917 were newly reported in the previous 7 days. The total deaths within 28 days of a positive test was 162,359 with 729 reported in the previous 7 days. Latest UK data is available from: https://coronavirus.data.gov.uk/
- As at 08/03/2022, WHO have reported 446,511,318 confirmed COVID-19 cases globally, with 1,172,188 reported in the previous 24 hours. There have been 6,004,421 deaths, of which 4,993 were reported in the previous 24 hours. Daily WHO situation updates are available from: <a href="https://covid19.who.int/">https://covid19.who.int/</a>

# Middle East respiratory syndrome coronavirus (MERS-CoV) - latest update from WHO and ECDC

- On 13/12/2021 WHO reported an additional case of Middle East Respiratory Syndrome coronavirus (MERS-CoV). Globally, 2,583 laboratory confirmed cases of human infection with MERS-CoV, including 888 associated deaths, have officially been reported to WHO since 2012.
   Source: WHO Global Alert and Response website: <a href="https://www.who.int/emergencies/disease-outbreak-news">https://www.who.int/emergencies/disease-outbreak-news</a>
- The majority of the MERS cases continue to be reported from the Middle East, and specifically from Saudi Arabia. Rapid risk assessments of the situation from ECDC, which contain epidemiological updates and advice for travellers and healthcare workers, are available from: <a href="https://ecdc.europa.eu/en/middle-east-respiratory-syndrome-coronavirus">https://ecdc.europa.eu/en/middle-east-respiratory-syndrome-coronavirus</a>
- Further updates and advice for healthcare workers and travellers are available from WHO: <a href="http://www.who.int/emergencies/mers-cov/en/">http://www.who.int/emergencies/mers-cov/en/</a> and from NaTHNaC: <a href="https://travelhealthpro.org.uk/news/237/mers-cov-update-travelhealthpro-country-pages">https://travelhealthpro.org.uk/news/237/mers-cov-update-travelhealthpro-country-pages</a>

# Human infection with avian influenza A(H7N9), China

- The latest WHO Influenza at Human-Animal Interface summary (14/12/2021 to 21/01/2022) reports that there have been no publicly available reports from China or other countries on influenza A(H7N9) in recent months. Since February 2013, a total of 1,568 laboratory-confirmed cases of human infection with avian influenza A(H7N9), including at least 616 deaths, have been reported:
   <a href="https://www.who.int/teams/global-influenza-programme/avian-influenza/monthly-risk-assessment-summary-http://www.fao.org/ag/againfo/programmes/en/empres/H7N9/Situation\_update.html">https://www.fao.org/ag/againfo/programmes/en/empres/H7N9/Situation\_update.html</a>
- The risk of international spread of avian influenza A(H7N9) is considered to be low at present. However, it is
  important that clinicians are aware of the possibility of human infection with animal influenza, in persons
  presenting with severe acute respiratory disease, while travelling or soon after returning from an area where
  avian influenza is a concern. WHO Global Alert & Response updates:
  <a href="https://www.who.int/emergencies/disease-outbreak-news">https://www.who.int/emergencies/disease-outbreak-news</a>

Links:

Public Health Wales influenza surveillance webpage:

http://www.wales.nhs.uk/sites3/page.cfm?orgid=457&pid=25480

Public Health Wales COVID-19 data dashboard:

https://public.tableau.com/profile/public.health.wales.health.protection#!/vizhome/RapidCOVID-19virology-

**Public/Headlinesummary** 

**GP Sentinel Surveillance of Infections Scheme:** 

http://www.wales.nhs.uk/sites3/page.cfm?orgid=457&pid=27918

NICE influenza antiviral usage guidance:

http://www.nice.org.uk/Guidance/TA158

Wales influenza information:

https://phw.nhs.wales/topics/flu/

**England influenza and COVID-19 surveillance:** 

https://www.gov.uk/government/statistics/national-flu-and-covid-19-surveillance-reports

Scotland seasonal respiratory surveillance:

https://beta.isdscotland.org/find-publications-and-data/population-health/covid-19/weekly-national-seasonal-respiratory-report/

Northern Ireland influenza surveillance:

https://www.publichealth.hscni.net/directorate-public-health/health-protection/seasonal-influenza

**European Centre for Communicable Disease:** 

http://ecdc.europa.eu/

European influenza information:

http://flunewseurope.org/

Advice on influenza immunisation (for NHS Wales users)

http://nww.immunisation.wales.nhs.uk/home

For further information on this report, please email Public Health Wales using: surveillance.requests@wales.nhs.uk