

## Adult Post-Transplant Immunisation Protocol (Month 24)

Time Post Transplant	Details
24 months	1. Measles, mumps, rubella (German measles) 1. One injection (MMR VaxPRO <sup>2</sup> ® or Priorix®)

### Notes:

1. The annual (seasonal) influenza vaccine is to be given every autumn.
2. Live vaccines are contra-indicated for at least 24 months post-transplant due to the risk of vaccine-induced illness.

**MMR is only to be given to patients in remission, without GvHD, not on immunosuppression and at least 8 months from the last immunoglobulin administration.**

**It is the responsibility of the transplant physician to determine (at the month 24 visit) if these conditions have been met and to send this form with a covering letter to the GP requesting that the MMR vaccine is given.**

3. **To the GP:** To **document compliance** with this document, please complete immunisation record (CLIN-FRM-066).
4. To **monitor efficacy** of this protocol, please assess vaccine antibody levels (where available) to all vaccines administered under this protocol annually to determine the need for boosters (see CLIN-FRM-006, CLIN-FRM-013 and CLIN-FRM-018). These tests will be requested by the transplant team.

### References:

1. Tomblyn M et al (2009), Guidelines for preventing infectious complications among hematopoietic cell transplantation recipients: a global perspective. Biol Blood Marrow Transplant 15:1143-1238.
2. ECIL-7 consensus document (2017), Guidelines for vaccination of patients with haematological malignancies and HSCT recipients.

SOP or Protocol reference:	N/A
Approved by: SWBMT Group	Date of approval: 26 January 2018
<b>It is the responsibility of the user of this document to confirm that this is the current version by checking on the Cardiff and Vale UHB intranet site</b>	